

Molecular Subtyping of *Salmonella enterica* serovar Enteritidis Isolated from Taiwan during 1992-1998 by Amplified Fragment Length Polymorphism and Pulsed-Field Gel Electrophoresis

TSUNG-YU TSAI¹, WEI-CHEN LUO¹, KUANG-LO CHEN² AND TZU-MING PAN^{1*}

¹ Institute of Microbiology and Biochemistry, National Taiwan University, 1, Sec. 4, Roosevelt Rd., Taipei City 106, Taiwan, R.O.C.

² Division of Laboratory Research and Development, Center for Disease Control, Taipei, Taiwan R.O.C.

(Received: April 1, 2005; Accepted: September 16, 2005)

ABSTRACT

Over the past 10 years, *Salmonella enterica* serovar Enteritidis has emerged as a new serovar and a frequent cause of food-borne diseases in Taiwan. We used two molecular subtyping methods to investigate the development of the infections by *S. enterica* Enteritidis in Taiwan. Twenty-eight strains of *S. enterica* Enteritidis isolated from Taiwan during the period of 1992 to 1998 were collected. The primer pairs, *MseI*+*C/EcoRI*+0 and *MseI*+*G/PstI*+*C*, were used in amplified fragment length polymorphism (AFLP) analyses, and the restriction enzyme, *AvrII*, was used in pulsed-field gel electrophoresis (PFGE). Twenty-two AFLP profiles with 53~59 fragments in each profile were found by using the primer pair, *MseI*+*G/PstI*+*C*. In the PFGE analysis, 7 PFGE types were identified. Some of the collected strains, especially the imported strains, had closer relationships with the SE 02 strain isolated from Taipei County. As our results indicate, AFLP analysis was time-saving, easy to perform, and highly discriminative. In this study, AFLP and PFGE were used to analyze the *S. enterica* Enteritidis isolated from patients in Taiwan. These results provide the epidemiological distribution of these isolated strains and additional evidence to illustrate the sources of the food-borne pathogens.

Key words: *Salmonella enterica* serovar Enteritidis, amplified fragment length polymorphism (AFLP), pulsed-field gel electrophoresis (PFGE)

INTRODUCTION

There are more than 200 diseases which are transmitted through food⁽¹⁾. The causes of food-borne illnesses include bacteria, viruses, parasites, toxins, and metals. In the USA, food-borne diseases cause an estimated of 6 to 81 million illnesses, and 9000 deaths each year⁽²⁻⁶⁾. Between 1986 and 1995, 852 outbreaks of food-borne disease which involved 26,173 cases and 20 deaths were reported in Taiwan; of these 852 reported outbreaks, 555 (65%) were caused by bacterial pathogens⁽⁷⁾.

Food poisoning caused by *Salmonella* is prevalent in many countries. *S. enterica* serovar Enteritidis is one of the most common serovars responsible for human illness, and is usually spread via contaminated egg or poultry meat. It can infect the ovary and the oviduct of poultry such that eggs can be contaminated and cause infections⁽⁸⁾. In Taiwan, outbreaks and sporadic infections caused by this pathogen were rare prior to 1994^(7,9); however, a dramatic increase in this type of infection was observed after 1995, and *S. enterica* Enteritidis caused 6 cases of infections between 1995 and 1997⁽¹⁰⁾.

Pulsed-field gel electrophoresis (PFGE) is one of the best molecular typing methods for analyzing the differences between closely related organisms, and is frequently applied to investigate the epidemiology of outbreaks. On the other hand, amplified fragment length polymorphism (AFLP) is a polymerase chain reaction-based method for rapid screening of genetic diversity⁽¹¹⁾. The latter is a cheap, easy, fast, and reliable method to generate hundreds of informative genetic markers⁽¹²⁾; moreover, it has been used for subtyping the *E. coli* strains including *E. coli* O157:H7^(13,14,15). We have collected the strains of *S. enterica* Enteritidis isolated from clinical patients reported to the Center for Disease Control (CDC) Taiwan, ROC from 1992 to 1998, and we used both AFLP and PFGE to investigate the relationships of *S. enterica* Enteritidis. Aside from the combination of the restriction enzymes, *MseI* and *EcoRI*, the *MseI* and *PstI* combination was also applied in order to create a more distinctive AFLP analysis condition. For the PFGE analysis, the restriction enzyme *AvrII* was used to determine the strains isolated. The results of these 2 molecular subtyping methods were analyzed and compared to illustrate the development of the infections by *S. enterica* Enteritidis in Taiwan.

* Author for correspondence. Tel: +886-2-33664519; Fax: +886-2-23627044; E-mail: tmpan@ntu.edu.tw

MATERIALS AND METHODS

I. Bacterial Strains

Twenty-eight isolates of *S. enterica* Enteritidis strains were used in this study. Twenty-seven of these strains were collected by the CDC, Taiwan and identified by biochemical and serotype tests. One strain was obtained from the Bureau of Food and Drug Analysis, Department of Health, Taiwan, ROC. All strains were isolated from outbreaks of food-borne gastroenteritis clinical stool samples or food samples from 1992 to 1998 in Taiwan. One strain, *Salmonella enteritidis* BCRC 10744 (purchased from the Bioresource Collection and Research Center, Food Industry Research and Development Institute, Hsinchu, Taiwan) was analyzed in this study as well. All strains were also re-identified by the multiplex PCR with the primers and conditions developed by our laboratory in previous study⁽¹⁶⁾.

II. AFLP Analysis

Genomic DNA was extracted using the Wizard[®] Genomic DNA purification kit (Promega Co., Madison, WI, USA), and the kit was evaluated according to the manufacturer's instructions. AFLP was performed and improved by methods described elsewhere⁽¹⁷⁻¹⁹⁾. In brief, the extracted DNA was subjected to restriction-ligation reactions for 3 hr in a total volume of 10 µL. This consisted of 10 ng of genomic DNA, 5 U of *EcoRI* or *PstI*, 1 U of *MseI*, 1 U of T4 DNA ligase (Epicentre, Madison, WI, USA), 1× ligase buffer (33 mM Tris-acetate pH 7.8, 66 mM potassium acetate, 10 mM magnesium acetate and 0.5 mM DTT) and 0.2 µg of each adaptor, as shown in Table 1. The restriction-ligation products were diluted 10 to 20-fold since the polymerase chain reaction (PCR) template for pre-amplified reaction was under the touch-down PCR conditions. It was then heated at 60°C for 30 min, and then stored at 4°C. The PCR products were diluted 10 to 20-fold as the selective PCR templates, and then the selective PCR was carried out with the primer pairs *MseI*+C/*EcoRI*+0,

MseI+G/*EcoRI*+G, *MseI*+G/*PstI*+G, and *MseI*+G/*PstI*+C which were screened in 50 different combinations. The selective extension nucleotide included adenine (A), thymine (T), cytosine (C), guanine(G) or non-extension (0) which extended on the end of the primer pairs. The reaction consisted of 200 µM dNTP, 0.2 µM for *MseI*+X selective primer, 0.04 µM for *EcoRI*+X or *PstI*+X selective primer (shown in Table 1), 1 U of *Taq* polymerase (TaKaRa *Taq*TM) and PCR reaction buffer [100 mM Tris-HCl (pH 8.3), 500 mM KCl, and 15 mM MgCl₂]. For selective amplified reactions, the PCR condition was 33 cycles at 94°C for 1 min, 60°C for 1 min and 72°C for 2.5 min. The PCR products were removed for capillary electrophoresis by sequence analyzer (3100, Applied Biosystems, Foster City, CA, USA) and Genescan TAMRA-500 (Applied Biosystems) as the internal standard in each sample⁽¹³⁾. The profiles of all strains were exported by Genescan[®] software (Applied Biosystems) and dendrograms for cluster analysis were performed by multivariate analysis of ecological data software, PC-ORD 4.0 (MJM software design, Glenenden Beach, OR, USA) using the Pearson correlation coefficient and the unweighted pair-group method using arithmetic average (UPGMA) algorithm.

III. PFGE Analysis

Pulsed-field gel electrophoresis was performed and based on the method developed by Barrett *et al.*⁽²⁰⁾. In brief, strains were grown in Trypticase Soy Broth (TSB) at 37°C to an optical density of 1.4~1.5 at 610 nm. Cells were washed twice and re-suspended in 75 mM NaCl / 25 mM EDTA. Bacterial cell suspension was carefully mixed with an equal volume of melted 1% chromosomal grade agarose (Bio-Rad Laboratories, Hercules, CA, USA), and the mixture was dispensed into 1.5-mm-thick block molds (Bio-Rad). After solidification, the plugs were transferred to microcentrifuge tubes containing lysis buffer (50 mM Tris pH 8.0, 50 mM EDTA pH 8.0, 1% N-laurylsarcosine, and 1 mg/mL proteinase K), and were then incubated overnight at 53°C. After then, the plugs were washed twice for 30 min in 10 nM Tris / 1 mM EDTA (pH 8.0) (TE) containing 1.5 mM phenylmethylsulfonyl fluoride (PMSF), and 4 times in TE buffer without PMSF. Two restriction enzymes, *XbaI* and *AvrII* (New England Biolabs, Beverly, MA, USA), were evaluated for typing these strains. The enzyme concentration, buffer, and incubation temperature were those recommended by the manufacturer for a 4-hr digestion of agarose-embedded DNA. Restriction fragments were separated by electrophoresis through 1% PFGE agarose (Bio-Rad Laboratories) in 0.5× Tris-borate-EDTA buffer at 14°C in a CHEF DR-II apparatus (Bio-Rad Laboratories) for 22 hr at a constant voltage of 6 V/cm. The pulse time was ramped from 2 sec in the beginning to 40 sec in the end. After PFGE, the gel was stained with ethidium bromide (0.2 g/mL) and photographed under UV transillumination by Kodak Electrophoresis Documentation and Analysis System 290 (Kodak Co., CA, USA).

Table 1. The oligonucleotides used in this study for the amplified fragment length polymorphism

| Name | Description | Sequence |
|------|----------------------|-----------------------------|
| MAD1 | <i>MseI</i> adaptor | 5'-GACGATGAGTCCTGAG-3' |
| MAD2 | | 3'-TACTCAGGACTC AT-5' |
| EAD1 | <i>EcoRI</i> adaptor | 5'-CTCGTAGACTGCGTACC-3' |
| EAD2 | | 3'-CTGACGCATGG TTAA-5'. |
| PAD1 | <i>PstI</i> adaptor | 5'-CTCGTAGACTGCGTACATGCA-3' |
| PAD2 | | 3'-CATCTGACGCATGT-5' |
| MSEP | <i>MseI</i> primer | 5'-GATGAGTCCTGAG TAA-3' |
| ECOP | <i>EcoRI</i> primer | 5'-GACTGCGTACC AATTN*-3' |
| PSTP | <i>PstI</i> primer | 5'-GACTGCGTACATGCAN*-3' |

*N: selective extension nucleotide including adenine (A), thymine (T), cytosine (C), guanine (G) or non-extension (0) which extended on the end of the primer pairs.




Figure 1. The profiles of the AFLP analysis results generated by Genescan[®] software by using the *MseI/EcoRI* condition. (A) SE 01; (B) SE 10; (C) BCRC10744.




Figure 2. The percentages of genetic similarity between 29 strains of *S. enterica* Enteritidis using the *MseI/EcoRI* condition as determined by multivariate analysis of ecological data software, PC-ORD 4.0.




Figure 3. The percentages of genetic similarity between 29 strains of *S. enterica* Enteritidis using the *MseI/PstI* condition as determined by multivariate analysis of ecological data software, PC-ORD 4.0.

Dendrograms for cluster analysis were performed by Bio-Profil[®] Image Analysis Software (Vilber Lourmat Co., Marne La Vallee, France) using the Pearson correlation coefficient and the UPGMA algorithm.

RESULTS AND DISCUSSION

AFLP analysis performed with 2 sets of enzyme combinations (*MseI/EcoRI* and *MseI/PstI*) yielded amplified fragments ranging from 60 to 600 bp in size (Shown in

Figure 1). Fifty different combinations needed to be screened, and the selective primer pair combinations, *MseI+C/EcoRI+0*, *MseI+G/EcoRI+G*, *MseI+G/PstI+G*, and *MseI+G/PstI+C* were utilized to type these strains. Under the condition of *MseI/EcoRI* (ME) enzyme combination, the primer pair *MseI+C/EcoRI+0* yielded 14 AFLP profiles (shown in Table 2) with 19~23 fragments in each profile. The selective bases of the primers were determined using *MseI/PstI* (MP) enzyme combination, and the primer pair *MseI+G/PstI+C* was found to be the most effective for the epidemiological typing of the 29 strains of *S. enterica*

Table 2. Subtypes of the type strain and 28 *Salmonella enterica* Enteritidis strains isolated during 1992-1998 by AFLP and PFGE

| Strains | Source | | AFLP | | PFGE |
|-------------------------------------------|--------|----------------------|------|-----|--------------|
| | Year | Location | ME | MP | <i>AvrII</i> |
| SE 01 | 1992 | Hualien City | E1 | P1 | A2 |
| SE 02 | 1992 | Taipei County | E2 | P2 | A1 |
| SE 03 | 1992 | Philippines imported | E2 | P3 | A1 |
| SE 04 | 1992 | Philippines imported | E2 | P3 | A1 |
| SE 05 | 1993 | Hualien City | E3 | P4 | A1 |
| SE 06 | 1993 | Hualien City | E4 | P5 | A1 |
| SE 07 | 1994 | Taipei City | E5 | P6 | A1 |
| SE 08 | 1995 | Taipei County | E6 | P3 | A3 |
| SE 09 | 1995 | Taipei County | E7 | P7 | A1 |
| SE 10 | 1995 | Taipei County | E8 | P8 | A1 |
| SE 11 | 1995 | Taipei County | E9 | P9 | A1 |
| SE 12 | 1995 | Taipei County | E10 | P3 | A1 |
| SE 13 | 1996 | Taipei County | E10 | P10 | A1 |
| SE 14 | 1996 | Taoyuan County | E11 | P3 | A1 |
| SE 15 | 1996 | Taipei City | E5 | P5 | A4 |
| SE 16 | 1996 | Taipei City | E2 | P11 | A1 |
| SE 17 | 1996 | Taoyuan County | E2 | P12 | A1 |
| SE 18 | 1997 | Taoyuan County | E2 | P13 | A1 |
| SE 19 | 1997 | Taoyuan County | E2 | P13 | A1 |
| SE 20 | 1997 | Taipei City | E2 | P14 | A1 |
| SE 21 | 1997 | Taipei County | E2 | P15 | A1 |
| SE 22 | 1997 | Taoyuan County | E2 | P16 | A1 |
| SE 23 | 1998 | Taipei County | E12 | P17 | A5 |
| SE 24 | 1998 | Taipei City | E13 | P18 | A6 |
| SE 25 | 1998 | Taoyuan County | E2 | P19 | A1 |
| SE 26 | 1998 | Taipei County | E2 | P20 | A1 |
| SE 27 | 1998 | Taipei County | E2 | P21 | A1 |
| SE 28 | 1998 | NR* | E2 | P21 | A1 |
| <i>S. enterica</i> Enteritidis BCRC 10744 | | | E14 | P22 | A7 |

*No report.

Enteritidis in AFLP analysis. Using this technique, 22 AFLP profiles (shown in Table 2), with 53~59 fragments in each profile, were determined. The percentages of genetic similarity between the 29 strains of *S. enterica* Enteritidis, using ME and MP conditions, are shown in Figures 2 and 3. Using ME, 11 strains had a unique profile (SE 01, 05, 06, 08, 09, 10, 11, 14, 23, 24, and BCRC10744) and there were 14 strains that shared AFLP profile E2 (Table 2). Strains SE 07 and SE 15, and SE12 and SE 13 shared AFLP profile E5 and E10 respectively. Using the MP results, 18 strains had unique profiles, and five strains (SE 03, 04, 08, 12, and 14) shared the same profile P3. Strains SE 06 and SE 15, SE 18 and SE 19, SE 27 and SE 28 shared AFLP profile P5, P13 and P21, respectively. The PFGE analysis of the 29 strains of *S. enterica* Enteritidis, using *AvrII* restriction enzyme, generated fragments ranging from 40 kb to 700 kb in size. The PFGE patterns are shown in Figure 4. Seven PFGE types were observed (Table 2), and the genetic diversity among these strains is illustrated by the dendrogram in

Figure 5. There were 23 strains that belonged to the PFGE type A1, and 6 strains which had unique types (A2 to A7).

Four strains (SE 09-12) from Taipei County, all from the same year, had different AFLP profiles. We compared the polymorphic fragments of these strains and found that the SE 11 strain was more different from the other strains isolated during the same time period. In the dendrogram, SE 09, SE 10, and SE 12 strains were closely related, and they were not related to the SE 11 strain. From the results of MP, it is evident that the strains SE 09, 10 and 12 have a higher relative relationship within the dendrogram, as shown in Figure 3. Although these 4 isolates had different AFLP profiles when using ME and MP, we hypothesized that the strains SE 09, 10, and 12 were more related based on the results of polymorphic fragments, dendrograms, and geographic location of infected cases. The case caused by strain SE 11 was a unique case of infection in Taipei County in 1995. On the other hand, the other 2 strains, SE 12 and 13, which were isolated from Taipei County




Figure 4. Pulsed-field gel electrophoresis patterns of *AvrII*-digested chromosomal DNA of SE 01~15 (A) and 16~28 (B) *S. enterica* Enteritidis isolates and type strain BCRC 10744. Lane M: lambda ladder marker; Lane 1~28: SE 01~SE 28; Lane 29: BCRC10744

in 1995 and 1996, had the same AFLP profiles under ME analysis. Although these 2 strains did not have the same profiles when using MP, and were of different time periods, the dendrogram of ME and the polymorphic fragments showed that these 2 strains were closely related. Therefore, we assume that these 2 isolates were closely related at that time. In summary, some of the 5 strains isolated from Taipei County at 1995 to 1996 were closely related, and




Figure 5. The percentages of genetic similarity between the 7 *AvrII* digestion pulsed-field gel electrophoresis patterns.

the infection may have had the same origin, especially the SE 12 and 13 strains from different years. The strains SE 16~19 isolated from Taipei City and Taoyuan County during 1995-1996 had a high degree of similarity in genotype and AFLP profiles when analyzed using ME and MP. Because of the close proximity of Taipei City and Taoyuan County, we can infer that these infected cases were connected with each other in those 2 years. Strain SE 01 had a unique PFGE type, A2, and strains SE 05 and 06, which were also isolated from Hualien, belonged to the A1 type. The results of PFGE were similar to those of AFLP with ME and MP, and indicated that strains SE 01, 05, and 06 were not related to each other. Based on these results, we can infer that these infected cases were not related to each other, even though they were all isolated in eastern Taiwan. Strain SE 27, isolated from Taipei County shared the same profile as strain SE 28 which was isolated from an unknown source of food sample, suggesting it may have originated from the same source as strain SE 27. When using both MP and ME for AFLP analysis, strain SE 02 showed close relatedness to the 2 imported strains from the Philippines (SE 03 and SE 04) isolated in the same year. Consequently, these 3 strains may have originated from the same source of imported infections.

Evidence from epidemiological and molecular studies of microorganisms suggests that strains are frequently transmitted across wide geographic distances⁽²¹⁾ due to the convenience of travel around the globe these days and increased human contacts. At the same time, agricultural, human, and food-borne diseases are dispersed easily and quickly and they are becoming increasingly difficult to prevent and control. As the investigation of Kim *et al.* demonstrated, the *Salmonella enterica* Serovar Typhimurium DT104 isolates were highly linked to the strains in Korea and United States because of animal and human circulation⁽²²⁾. On the other hand, the strains always have unique molecular profiles in the neighboring areas. In our previous study, the *E. coli* O157:H7 strains isolated in Taiwan have been investigated and were closely related

to each other, but significantly different from other foreign strains⁽²³⁾. It is important to establish molecular subtyping methods in order to trace and control the infection of foodborne pathogens. As the results of this study indicate, the infection of *S. enterica* Enteritidis has high geographic relationship which was centralized in Taipei City and Taoyan County, especially strains SE 09~13 which caused 4 outbreaks in 1995 and 1996. Strains SE 16~19 were in a similar situation from 1996 to 1997 in Taipei City and Taoyuan County. The closely related strains were isolated in the immediate areas during continuous years indicating that the dominant and highly related strains were circulating in these areas at northern Taiwan.

Molecular epidemiological typing of microbial isolates through the characterization of proteins or nucleic acids has been successfully applied to epidemiologic investigations of outbreaks of food-borne diseases^(24,25). During the past decade, some phenotyping methods were carried out to illustrate the relationship between pathogens, such as the phage type. Plasmid profile analysis and plasmid restriction fragment length polymorphism have been applied in the typing of *E. coli* O157⁽²⁶⁾. Several other molecular typing methods for analyzing the bacterial genomic DNA by using restriction enzymes have been developed in recent years. These methods include RFLP, PFGE, and AFLP, which all focused on analyzing the genomic DNA restrictive maps for epidemiologic investigation of infectious diseases^(20,27-33). Unfortunately, there has been no perfect method for the molecular typing of infectious pathogens; consequently the epidemiologists must carry out two or more molecular subtyping methods to identify the type of a strain. For example, strains SE 01, 05, and 06 were isolated from the same area, and the results from PFGE using *Xba*I indicated that these 3 strains had the same genotype, but the PFGE using *Avr*II and AFLP with ME and MP inferred that these 3 strains each had unique profiles. Therefore, it is necessary to combine and compare two or more results of molecular subtyping in epidemiology.

According to Struelens⁽³⁴⁾, different bacteria strains, different operators and procedural modifications can affect the results of AFLP. Desai *et al.* found that the different PCR condition was effective for genotyping *S. enterica* Enteritidis^(8,35). Lindstedt *et al.*⁽³⁶⁾ used the same primer pair, *Mse*I+C/*Eco*RI+0, for the AFLP analysis of *S. enterica* Enteritidis and the results showed that AFLP is a versatile molecular subtyping method. A good molecular typing method should be versatile, which means the method should be able to type any kind of bacterial strain. The situation they described could be solved by choosing different restriction enzymes or selective bases in the primer. The choice of restriction enzymes and the number of selective bases are critical in AFLP analysis⁽³⁷⁾. A highly frequent cutter enzyme will generate results which are too complex, while an extremely rare cutter enzyme will generate too few fragments. The former results are difficult to interpret while the latter ones may reduce the possibility of detecting polymorphism. At the same time, according to Aarts

et al.⁽³⁸⁾, 200 fragments are considered to hamper the interpretation; thus, most studies set the range between 50 to 200 fragments. However, different materials (bacteria, plant or animal cells) and a modification of procedures can still affect the range. In this study, after the restriction and selection it was found that ME has less fragments than MP and that the discriminatory power decreased appreciably.

Although AFLP with MP could differentiate *S. enterica* Enteritidis better than with ME in this study, strain SE 08, which has a unique profile when using PFGE with *Avr*II and AFLP with ME, could not be differentiated from other strains in AFLP with MP. Such results have also been reported from other studies^(39,40). Nair *et al.* hypothesized that the differences between the ability of PFGE and AFLP to differentiate strains may arise from the fact that the strain had no polymorphisms within the *Mse*I and *Eco*RI restriction sites in the sequences adjacent to the restriction sites. These sites are complementary to the selective base of the primer, as compared to the polymorphisms within the *Avr*II and *Xba*I sites (detected by PFGE)⁽⁴⁰⁾.

Although, at least theoretically, AFLP provides better differentiation of strains compared to PFGE, many researchers have considered the fact that an outbreak genotype should be identified on the basis of a combination of AFLP analysis and epidemiological context^(8,34). There already exists a system for standardizing the interpretation of PFGE patterns⁽⁴¹⁾, and a Pulsed-Net internet system was established to collect epidemiological data from all over the world so that inter-laboratory comparisons can be made⁽⁴²⁾. Also, the pattern interpretation for AFLP analysis needs to be standardized. To establish two or more molecular subtyping methods is becoming increasingly important for effective infection and disease control. In Taiwan, there is not any one department which has the all-inclusive capabilities to analyze food-borne pathogens for molecular subtyping; in addition, there is not any one molecular subtyping method that has been agreed upon as the official method to investigate food-borne or infectious diseases. In Taiwan, this present study is the first to propose regular methods of molecular subtyping for *S. enterica* Enteritidis infection, and for comparing the two kinds of subtyping methods. At the same time, there is no international organization set up to participate in this network which is so urgently needed right now. Therefore, it remains difficult to trace the source of any food-borne pathogens quickly, especially if that infection comes from a different country. The establishment and joining of a global network database is an important task if we wish to control the agricultural and food-borne pathogens, and reduce the losses resulting from the infection and spread of these pathogens.

CONCLUSIONS

In summary, we used the AFLP and PFGE methods to investigate the relationships between 28 isolates from Taiwan during 1992-1998. It was found that some of

these cases had a higher level of relationship, especially the imported strains, SE 03 and 04 which had the same genotype patterns as strain SE 02 isolated from Taipei County. The cases of infection by *S. enterica* Enteritidis in Taiwan was first determined in 1992, and had increased after 1995. These infections were centralized in northern Taiwan, specifically in Taipei City, Taipei County, and Taoyan County, where all had a large population. Our results showed that AFLP was efficient, easy to use and highly discriminative for a molecular subtyping method. It seemed to provide the most useful epidemiological information about *S. enterica* Enteritidis. To combine and compare two or more molecular subtyping information could provide more evidence to illustrate the sources of the food-borne pathogens.

ACKNOWLEDGEMENTS

We thank the Center for Diseases Control and the Bureau of Food and Drug Analysis, Department of Health, Taiwan, ROC for kindly providing the strains for this study.

REFERANCES

1. Byyan, F. L. 1982. Diseases Transmitted by Foods. Centers for Disease Control. Atlanta, U. S. A.
2. Ames, I. A. 1994. Foodborne Pathogens: Risks and Consequences. Council of Agricultural Science and Technology.
3. Archer, D. L. and Kvenberg, J. E. 1985. Incidence and cost of food-borne diarrheal disease in the United States. *J. Food Protect.* 48: 887-894.
4. Bennett, J., Holmberg, S., Rogers, M. and Solomon, S. 1987. Infectious and parasitic diseases. In "Closing the Gap: the Burden of Unnecessary Illness" pp. 102-214. Amler, R. and Dull, H. eds. Oxford Univ Press. New York, U. S. A.
5. Mead, P. S. and Griffin, P. M. 1998. *Escherichia coli* O157:H7. *Lancet* 352: 1207-1212.
6. Todd, E. C. D. 1989. Preliminary estimates of costs of food-borne disease in the United States. *J. Food Protect.* 52: 595-601.
7. Pan, T. M., Wang, T. K., Lee, C. L., Chien, S. W. and Horng, C. B. 1997. Food-borne disease outbreaks due to bacteria in Taiwan, 1986 to 1995. *J. Clin. Microbiol.* 5: 1260-1262.
8. Desai, M., Threlfall, E. J. and Stanley, J. 2001. Fluorescent amplified-fragment length polymorphism subtyping of the *Salmonella enterica* serovar Enteritidis phage type 4 clone complex. *J. Clin. Microbiol.* 39: 201-206.
9. Pan, T. M., Lin, C. S., Wang, T. K., Tsai, J. L., Ho, S. I. and Lee, C. L. 1998. Molecular subtyping of *Salmonella enterica* serovar *paratyphi* A from southeast Asia. *J. Food Drug Anal.* 6: 573-578.
10. Wang, T. K., Tsai, C. L., Lin, C. S. and Pan, T. M. 1999. Recent trend of salmonella food poisoning and serovars. *Epidemiology Bull.* 15: 1-6.
11. Mueller, U. G. and Wolfenbarger, L. L. 1999. AFLP genotyping and fingerprinting. *Trends Ecol. Evol.* 14: 389-394.
12. Russell, J. R., Fuller, J. D., Macaulay, M., Hatz, B. G., Jahoor, A., Powell, W. and Waugh, R. 1997. Direct comparison of levels of genetic variation among barley accessions detected by RFLPs, AFLPs, SSRs and RAPDs. *Theor. Appl. Genet.* 95: 714-722.
13. Arnold, C., Metherell, L., Clewley, J. P. and Stanley, J. 1999. Predictive modeling of fluorescent AFLP: A new approach to the molecular epidemiology of *E. coli*. *Res. Microbiol.* 150: 33-44.
14. Horvath, R., Dendis, M., Schlegelova, J., Ruzicka, F. and Benedik, J. 2003. A combined AFLP-multiplex PCR assay for molecular typing of *Escherichia coli* strains using variable bacterial interspersed mosaic elements. *Epidemiol. Infect.* 132: 61-65.
15. Zhao, S., Mitchell, S. E., Meng, J., Kresovich, S., Doyle, M. P., Dean, R. E., Casa, A. M. and Weller, J. W. 2000. Genomic typing of *Escherichia coli* O157:H7 by semi-automated fluorescent AFLP analysis. *Microbe. Infect.* 2: 107-113.
16. Pan, T. M. and Liu, Y. J. 2002. Identification of *Salmonella enteritidis* isolates by multiplex polymerase chain reaction. *J. Microbiol. Immunol. Infect.* 35: 147-151.
17. Arnold, C. L., Metherell, L. and Willshaw, G. 1999. Predictive fluorescent amplified-fragment length polymorphism analysis of *Escherichia coli*: High-resolution typing method with phylogenetic significance. *J. Clin. Microbiol.* 37: 1274-1279.
18. Gibson, J. R., Slater, E., Xerry, J., Tompkins, D. S. and Owen, R. J. 1998. Use of an amplified-fragment length polymorphism technique to fingerprint and differentiate isolates of *Helicobacter pylori*. *J. Clin. Microbiol.* 36: 2580-2585.
19. Vos, P., Hogers, R. and Bleeker, M. 1995. AFLP: A new technique for DNA fingerprinting. *Nucleic Acids Res.* 21: 4407-4414.
20. Barrett, T. J. 1997. Molecular fingerprinting of food-borne pathogenic bacteria: An introduction to methods, uses and problems. In "Food Microbiological Analysis: New Technologies". pp. 249-264. Tortorello, M. L. and Gendel, S. M. eds. Marcel Dekker. New York, U. S. A.
21. Davis, M. A., Hancock, D. D., Besser, T. E., Rice, D. H., Hovde, C. J., Digiacomio, R., Samadpour, M. and Call, D. R. 2003. Correlation between geographic distance and genetic similarity in an international collection of bovine faecal *Escherichia coli* O157:H7 isolates. *Epidemiol. Infect.* 131: 923-930.
22. Kim, S., Chun, S. G., Lim, O. Y., Park, M. S., Kang, Y. H., Park, Y. H. and Lee, B. K. 2004. Genomic relationship of *Salmonella enterica* Serovar Typhimurium DT104 isolates from Korea and the United States. *J.*

- Microbiol. 42: 14-19.
23. Tsai, T. Y., Luo, W. C., Wu, F. T. and Pan, T. M. 2005. Molecular subtyping for *Escherichia coli* O157:H7 isolated in Taiwan. Microbiol. Immunol. 49: 579-588.
 24. Holmberg, S. D., Wachsmuth, I. K., Hickman-Brenner, F. W. and Cohen, M. L. 1984. Comparison of plasmid profile analysis, phage typing, and antimicrobial susceptibility testing in characterizing *Salmonella typhimurium* isolates from outbreaks. J. Clin. Microbiol. 19: 100-104.
 25. Holmberg, S. D. and Wachsmuth, K. 1989. Plasmid and chromosomal DNA analyses in the epidemiology of bacterial diseases. In "Nucleic Acid and Monoclonal Antibody Probes: Applications in Diagnostic Microbiology". pp. 105-129. Swaminathan, B. and Prakash, G. eds. Marcel Dekker. New York, U. S. A.
 26. Swaminathan, B. and Matar, G. M. 1993. Molecular typing methods. In "Diagnostic Molecular Microbiology". pp. 26-50. Persing, D. H., Smith, T. F., Tenover, F. C. and White, T. J. eds. American Society for Microbiology. Washington, DC, U. S. A.
 27. Ackers, M. L., Mahon, B. E., Leahy, E., Goode, B., Damrow, T., Hayes, P. S., Bibb, W. F., Rice, D. H., Barrett, T. J., Hutwagner, L., Griffin, P. M. and Slutsker, L. 1998. An outbreak of *Escherichia coli* O157:H7 infections associated with leaf lettuce consumption. J. Infect. Dis. 177: 1588-1593.
 28. Graves, L. M., Swaminathan, B. and Hunter, S. B. 1999. Subtyping *Listeria monocytogenes*. In "Listeria, Listeriosis and Food Safety". pp. 279-298. Ryser, E. M. and Marth, E. H. eds. Marcel Dekker. New York, U. S. A.
 29. Jimenez, A., Barros-Velazquez, J., Rodriguez, J. and Villa, T. G. 1997. Restriction endonuclease analysis, DNA relatedness and phenotypic characterization of *Campylobacter jejuni* and *Campylobacter coli* isolates involved in food-borne disease. J. Appl. Microbiol. 82: 713-721.
 30. Maslanka, S. E., Kerr, J. G., Williams, G., Barbaree, J. M., Carson, L. A., Miller, J. M. and Swaminathan, B. 1999. Molecular subtyping of *Clostridium perfringens* by pulsed-field gel electrophoresis to facilitate food-borne-disease outbreak investigations. J. Clin. Microbiol. 37: 2209-2214.
 31. Threlfall, E. J., Hampton, M. D., Ward, L. R. and Rowe, B. 1996. Application of pulsed-field gel electrophoresis to an international outbreak of *Salmonella agona*. Emerg. Infect. Dis. 2: 130-132.
 32. Threlfall, E. J., Ward, L. R., Hampton, M. D., Ridley, A. M., Rowe, B., Roberts, D., Gilbert, R. J., van Someren, P., Wall, P. G. and Grimont, P. 1998. Molecular fingerprinting defines a strain of *Salmonella enterica* serotype Anatum responsible for an international outbreak associated with formula-dried milk. Epidemiol. Infect. 121: 289-293.
 33. Wachsmuth, K. 1986. Molecular epidemiology of bacteria infections: Examples of methodology and of investigations of outbreaks. Rev. Infect. Dis. 8: 682-692.
 34. Struelens, M. J. 1998. Molecular epidemiologic typing systems of bacterial pathogens: Current issues and perspectives. Mem. Inst. Oswaldo Cruz. 93: 581-585.
 35. Desai, M., Tanna, A. and Efstratiou, A. 1998. Extensive genetic diversity among clinical isolates of *Streptococcus pyogenes* serotype M5. Microbiology 144: 629-637.
 36. Lindstedt, B. A., Heir, E. and Vardund, T. 2000. Fluorescent amplified-fragment length polymorphism genotyping of *Salmonella enterica* subsp. Enterica serovar and comparison with pulsed-field gel electrophoresis typing. J. Clin. Microbiol. 38: 1623-1627.
 37. Ridiut, C. J. and Donini, P. 1999. Use of AFLP in cereals research. Trends Plant Sci. 4: 76-79.
 38. Aarts, H. J. M., van Lith, L. A. J. T. and Keijer, J. 1998. High genotyping of *Salmonella* strains by AFLP fingerprinting. Letters Appl. Microbiol. 26: 131-135.
 39. Møler, K., Jensen, T. K. and Bote, M. 1999. Amplified fragment length polymorphism and pulsed-field gel electrophoresis for subspecies differentiation of *Serpulina pilosicoli*. Anaerobe 5: 313-315.
 40. Nair, S., Schreiber, E. and Thong, K. L. 2000. Genotypic characterization of *Salmonella typhi* by amplified fragment length polymorphism fingerprinting provides increased discrimination as compared to pulsed-field gel electrophoresis and ribotyping. J. Microbiol. Methods 41: 35-43.
 41. Tenover, F. C., Arbeit, R. D. and Goering, R. V. 1995. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: Criteria for bacteria strain typing. J. Clin. Microbiol. 33: 2233-2239.
 42. Swaminathan, B., Barrett, T. J. and Hunter, S. B. 2001. PulseNet: The molecular subtyping network for foodborne bacterial disease surveillance, United States. Emerg. Infect. Dis. 7: 382-389.