

Molecular Authentication of Chinese Herbal Materials

YAN-BO ZHANG¹, PANG-CHUI SHAW², CHO-WING SZE¹, ZHENG-TAO WANG³ AND YAO TONG^{1*}

¹ School of Chinese Medicine, The University of Hong Kong, 10 Sassoon Rd., Pokfulam, Hong Kong Special Administrative Region, P.R. China

² Department of Biochemistry, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong Special Administrative Region, P.R. China

³ Department of Pharmacognosy, Chinese Pharmaceutical University, Jixiangnan Nanjing, P.R. China

(Received: May 19, 2006; Accepted: October 4, 2006)

ABSTRACT

Traditional Chinese medicine (TCM) has been playing a major role in health care in China for millennia. Accurate authentication is always necessary to prevent the target herbs from intentional and inadvertent adulteration with other plant species. Morphological and histological authentication is now commonly practiced but they are not precise enough to authenticate those herbs which are possibly substituted or adulterated by plants with similar shapes and tissue constructs. Ordinary chemical authentication was also introduced to TCM but it is often not reliable enough to produce easy-to-interpret results. Therefore, it is necessary to develop a more effective, accurate, reliable and sensitive technology for the authentication of herbs. DNA manipulation techniques developed for molecular biotechnology have been adapted to the authentication of herbs in recent years. These techniques comprise of the molecular markers, sequencing of specific genes, and sophisticated hybridization setups such as DNA microarrays. Underside, we review the development of current techniques for authentication. In addition, we also describe the use of molecular markers in authentication of the most studied Chinese herbs.

Key words: molecular authentication, rDNA, Chinese herbal materials

INTRODUCTION

For thousands of years, traditional Chinese medicine (TCM) has been practiced and has played a crucial role in the health care and in helping the Chinese nation flourish. In spite of the great advances of modern medicine, TCM is still the primary form of healing methods for many people in Asia⁽¹⁾. With its multi-target effects, TCM is particularly suitable for the treatment of modern diseases such as cardiovascular disease, asthma, and other long-term illnesses⁽²⁾. Furthermore, an increasing variety of healthy care products has been developed from TCM to meet the contemporary trend for "back to nature". Many famous multi-national medicine companies are now developing TCM jointly with Chinese companies. The World Health Organization has also been keen to pursue the development of TCM. The standardization and modernization of TCM depend on the authentication of the identity of Chinese medicinal materials. Therefore, the authentication and quality control have been the key for TCM to enter the world market⁽³⁾.

Up till now, traditional morphological inspection is still widely used to distinguish the herbs. Morphological approach includes the inspection of shape, color, texture and odor of the herbs. For example, in the traditional authentication of *Radix Codonopsis*, the following criteria are used: the root is long cylindrical, slightly curved, 10-35 cm long and 0.4-2 cm in diameter⁽⁴⁾ and the odour is characteristic, aromatic and tastes sweet⁽⁵⁾. The morpho-

logical inspection to authenticate TCM is simple and direct but its accuracy depends heavily on the examiners, which are sometimes subjective.

Histological techniques based on microscopic examinations are used to reveal the characteristics of tissue structure and arrangement in cork cell, cortex, sieve tubes, xylem vessels and cell components or content of a manufactured product. The thickness of the exodermal cell walls, diameter of exodermis, number of transfusion cells, and number of vascular bundles were used as markers for identifying the plants source of *Herba Dendrobii*⁽⁶⁾. The transverse section of *Radix Codonopsis* has over ten rows of cork cells. There are stone cells present at the outer side. The cortex is narrow⁽⁷⁾. Histological identification is not applicable to modern herbal drugs, for example, herbal capsules, troche and pills. Related species may share similar histological characteristics, making this approach not so accurate.

Chemical authentication emphasizes on the analysis of chemical constituents. Characteristic compositions are used for the differentiation. Thin layer chromatography (TLC) is the most common techniques to assess the chemical constituents of medicinal materials. For examples, TLC was used to identify *Tribulus terrestris*⁽⁸⁾ and *Fructus Xanthii*⁽⁹⁾. High performance liquid chromatography (HPLC) has become the routine procedures to identify herbal materials. For example, HPLC has been used to analyze the chemical profiles of cassia bark (cortex *cinnamomi*)⁽¹⁰⁾, to generate fingerprints for *Psoralea corylifolia*⁽¹¹⁾, to authenticate *Ephedra*⁽¹²⁾ and to evaluate the quality of *Radix Salviae Miltiorrhizae*⁽¹³⁾. An accurate

* Author for correspondence. Tel: +852-25890436; Fax: +852-28725476; E-mail: tongyao@hku.hk

and reproducible reversed-phase HPLC was developed to determinate the content of atractylenoide III, which is known as the active constituent of *Codonopsis pilosula*⁽¹⁴⁾. Besides HPLC, other chemical approaches have also been developed including ultraviolet spectroscopy⁽¹⁵⁾, infrared spectroscopy⁽¹⁶⁾, gas chromatography/mass spectrometry⁽¹⁷⁾, liquid chromatography/mass spectrometry⁽¹⁸⁾ and liquid chromatography/mass spectrometry/mass spectrometry⁽¹⁹⁾. The compositions and relative amount of chemicals in a species may have also been developed with the growing conditions, harvesting periods, post-harvest processes and storage. The variation of chemical compositions may hinder the authentication, and in some instances, this can be misleading if the samples are deliberately adulterated with a marker compound. Moreover, it is difficult to distinguish closely related species due to similar chemical compounds.

I. Molecular Markers for the Authentication of TCM

(I) DNA-based Molecular Markers

DNA-based markers have now become a popular means for the identification and authentication of TCM from plants and animals. Major techniques include random amplified polymorphic DNA (RAPD)⁽²⁰⁻²³⁾ and arbitrarily-primed polymerase chains reaction (AP-PCR)⁽²⁴⁻²⁵⁾, polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP)⁽²⁶⁻²⁸⁾, amplified fragment length polymorphism (AFLP)⁽²⁹⁾, direct amplification of length polymorphism (DALP)⁽³⁰⁾, specific sequence characterized amplified region (SCAR)⁽³¹⁻³²⁾ and short sequence repeat (SSR)⁽³³⁾. Among these, RAPD technique was applied early to differentiate *C. pilosula* species from different areas in China. DNA fingerprints were used to distinguish Chinese herb Dangshen, the root of *C. pilosula* from different localities in China. This method may be applicable to locality authentication of other Chinese herbal materials (CHM)⁽³⁴⁾. The phylogenic trees for *C. lanceolata* have been constructed for phylogenic analysis⁽³⁵⁾. Genetic diversity, relationship and molecular authentication of total 8 wild populations of *Dendrobium officinale* were investigated using RAPD markers. Distinct genetic differences and extensive genetic diversity were presented among the wild populations. RAPD markers are informative and useful tools for the evaluation and authentication of wild populations of *D. officinale*⁽³⁶⁾. Recently, RAPD was combined with other methods for the identification of Chinese medicines. For example, RAPD and Eastern blotting analyses using ginsenoside Rb1 and Rg1 monoclonal antibodies were employed to identify *Panax notoginseng*, *P. quinquefolius* and *P. japonicus*. RAPD was first used to differentiate the species of *Panax* spp. and thus the absence of ginsenoside Rc in the extract of *P. notoginseng* in the Eastern blot confirmed the identity of this species⁽³⁷⁾. *Fritillaria pallidiflora* is a commonly used antitussive herb. The differentiation of eight *F.*

pallidiflora species is limited by the current morphology-based and chemical methods. Therefore diagnostic PCR and PCR-RFLP have been established to differentiate *Fritillaria* species⁽³⁸⁾. In Brazil, *Plectranthus* species are known as “boldo” and have been commonly used for analgesic and dyspeptic purposes. *Plectranthus* spp. need to be well identified in order to be used commercially and AFLP DNA patterns have been used to distinguish different *Plectranthus* species⁽³⁹⁾.

(II) DNA Sequencing-based Markers

DNA polymorphisms were studied by determining the nucleotide sequence in a defined region and aligning the sequence with homologous regions of related organisms⁽⁴⁰⁾. This approach provides a highly reproducible and informative analysis and can be adapted to various levels of discriminatory potential by choosing appropriate regions of the genome. Currently, DNA sequencing is applied to distinguish species and study phylogenetic relationship, population genetics, systematics and evolution⁽⁴¹⁾. There are many reports concerning the application of DNA sequence-based markers to differentiate TCM from its substitutes or adulterants. Most of them involves the sequencing of internal transcribed spacer (ITS) ribosomal DNA (rDNA)⁽⁴²⁻⁴⁴⁾, 5S rDNA gene⁽⁴⁵⁻⁴⁶⁾, 18S rDNA and *trnK* genes⁽⁴⁷⁾, cytochrome b⁽⁴⁸⁾ and chloroplast DNA (cpDNA)⁽⁴⁹⁾. First, the ITS rDNA region has become an important gene locus for the molecular systematic investigation of angiosperms at the interspecific and intraspecific levels. Specific PCR primers are positioned on the conserved rDNA genes (18S, 5.8S, 28S) to amplify the entire ITS spacer region (Figure 1). The ITS region of rDNA, defined as the unit containing the ITS1 spacer, 5.8S rDNA gene and ITS2 spacer, has been proven to be a useful gene for screening different species of TCM. ITS rDNA region is unable to differentiate *Codonopsis* species because the sequences are highly conserved⁽⁵⁰⁾. On the other hand, the physical maps of chloroplast DNA (cpDNA) of *Codonopsis* genus were constructed and they are corresponding to the pollen morphology. It was suggested that cpDNA gene order mutations make an excellent phyloge-




Figure 1. Schematic diagram of the nuclear rDNA internal transcribed spacer region. The three rDNA subunits: 18S, 5.8S and 28S are separated by internal transcribed spacers (ITS1 and ITS2). Arrows indicate the annealing sites of the primers used for PCR amplification. 18b: 5'-TAG AGG AGG GAG AAG TCG TA-3'; 18d: 5'-CAC ACC GCC CGT CGC TCC TAC CGA-3'; 28cc: 5'-ACT CGC CGT TAC TAG GGG AA-3'; 5.8b: 5'-TGA AGA ACG TAG CGA AAT GCG-3'; 5.8d: 5'-AAC CAT CGA GTC TTT GAA CGC A-3'.

netic marker⁽⁵¹⁾. Several laboratories have also employed ITS rDNA region for the authentication of *Dendrobium* species and its product Shihu. The sequences of ITS2 regions of 16 *Dendrobium* species differ from one another by an average of 12.4% and differ from non-orchids and *Pholidota* (an adulterant of *Dendrobium*) by 29.8% and 18.8%, respectively⁽⁵²⁾. The ITS2 regions could be thus adopted as a molecular marker for differentiating medicinal *Dendrobium* species from one another and also from non-orchids and adulterants. The ITS sequences were also used to analyze a class of Herba Dendrobii with thin yellowish stems known as Huangcao Shihu⁽⁵³⁾. There are two nucleotide differences in the ITS region between F type (can be processed to Fengdou Shihu) and H type (cannot be processed to Fengdou Shihu) of *D. officinale* in China⁽⁵⁴⁾. The ITS region was used to authenticate Fengdou Shihu⁽⁵⁵⁾ and to distinguish *D. chrysanthum* from its relative species⁽⁵⁶⁾. Further, the whole ITS1-5.8S-ITS2 rDNA regions of 28 *Dendrobium* species were sequenced (GenBank accession number AY485692-AY485719). The average difference of the ITS1 is 34.62% between *Dendrobium* and non-orchids, and 22.31% between the *Dendrobium* and the orchids; the interspecific difference among the *Dendrobium* species is 13.14% (Table 1), indicating that ITS1 may also be used to differentiate the concerned *Dendrobium* species⁽⁵⁷⁾. Secondly, 5S rDNA gene of CHM was also explored. Radix Adenophorae (Shashen) is derived from the roots of *Adenophora stricta* and *A. tetraphylla*. Twelve species and varieties of *Adenophora* and *Glehnia*, however, have been used as substitutes or adulterants of Radix Adenophorae in the South East Asia markets. The 5S rDNA spacer domains (approximately 250 bp) were amplified by PCR from genomic DNAs from *A. stricta*, *A. tetraphylla*, *A. hunanensis* and *G. littoralis*, and sequences. The diversity in DNA sequences and restriction enzyme mapping among various species were found, which could serve as markers for the authentication of Radix Adenophorae⁽⁵⁸⁾. *Rhizoma Curcumae* (Ezhu) has been used to remove blood stasis and to alleviate pain for centuries. The 5S rDNA spacer domains of five *Curcuma* species, including the common adulterants of this herb, were amplified and sequenced. The diversity in DNA sequenced was used for the quality control of these *Curcuma* species⁽⁵⁹⁾. Thirdly, 18S, *trnK*, 12S, cytochrome b genes were also investigated. The six botanical origins of Chinese and Japanese *Curcuma* drugs were determined based on the comparison of 18S rDNA gene and *trnK* gene sequences. To develop a more convenient identification method, amplification-refractory mutation system (ARMS) analysis of both gene regions was performed. By the ARMS method, and the information on the region of production, the identification of *Curcuma* plants was achieved. The ARMS method for the *trnK* gene was also useful for authentication of *Curcuma* drugs⁽⁶⁰⁾. Chloroplast *trnK* gene and nuclear 18S rDNA sequences of 13 *Panax* taxa, collected mainly from Sino-Japanese floristic regions, were investigated in order to construct the phylo-

Table 1. The average percentage differences of the ITS1 rDNA gene among varies *Dendrobium*, orchids and non-orchids samples

Samples	Maximum (%)	Average (%)	Minimum (%)
Intraspecific	2	1.03	0
Interspecific	25	13.14	5
Other orchids	34	22.31	16
Non-orchids	43	34.62	31

genetic relationship and to assist the taxonomic delimitation within this genus⁽⁶¹⁾. To distinguish the Chinese crude drug Sailonggu (bone of plateau zokor, *Myospalax baileyi*) from its substitutes, two pairs of allele-specific diagnostic primers (SL1L/SL1H and SL2L/SL2H) were designed based on the mitochondrial 12S rDNA and cytochrome b genes sequences of the bamboo rat (*Rhizomys sinensis*) and black lipped pika (*Ochotona curzoniae*). Each of the two diagnostic primer pairs can be used to distinguish crude drug Sailonggu from its substitutes or adulterants. In addition, the results of sequence alignment and phylogenetic analysis are consistent with that of the relative-specific diagnostic PCR analysis⁽⁶²⁾. Two regions inside the chloroplast *trnK* were selected for the authentication of *Atractylodes* Rhizome (Byaku-jutsu) and *A. Lancea* Rhizome (So-jutsu). By comparing the nucleotide sequence data sets, it is possible to discriminate Byaku-jutsu and So-jutsu and also to identify the original plant species of each crude drug specimen⁽⁶³⁾. Last, Chloroplast chlB gene encoding the subunit B of light-independent protochlorophyllide reductase was amplified from herbarium and crude drug specimens of *Ephedra sinica*, *E. intermedia*, *E. equisetina*, and *E. przewalskii*, for the authentication of the corresponding crude drugs obtained in the Chinese market⁽⁶⁴⁾.

(III) DNA Microarray-based Makers

At present, high-density miniaturized microarrays (Biochip) have emerged as promising tools for the high throughput analysis of genomic data. DNA microarray revolutionizes the traditional way of one gene per experiment for the genome studies⁽⁶⁵⁾. Armed with the ITS sequences, microarray of the ITS1-5.8S-ITS2 regions from 28 *Dendrobium* species, two other orchids and two non-orchids were generated. Distinctive hybridization profile showed that 24 *Dendrobium* species may be differentiated from one another. The differentiation of *D. officinale* and *D. hercoglossum*, *D. nobile* and *D. moniliforme* was achieved by 5S rDNA array. This work has shown that ITS microarray could be used not only to establish the identities of the various *Dendrobium* species, but also to authenticate the medicinal *Dendrobium* from adulterant orchids⁽⁵⁴⁾. To develop a rapid, accurate and sensitive method for identifying the source plant from the product, ITS microarray was employed to authenticate the Herba Dendrobii from two medicinal formulations. In

this experiment, the ITS1-5.8S-ITS2 sequences were used as probes and the ITS2 sequences as target. The Herba Dendrobii in formulation A, which contained nine herbal materials, was found to be *D. nobile* (Figure 2). On the other hand, formulation B, which contains 12 components, was tested and the Herba Dendrobii in this formulation was found to be *D. lohohense*. The latter species in fact is not listed in the Chinese Pharmacopoeia and hence is a substitute of Herba Dendrobii⁽⁵⁵⁾. These studies provide the very examples of DNA microarray technology in tracing a medicinal component from complex medicinal mixture. In addition, microarray technology has also been used to authenticate ginseng⁽⁶⁶⁾ and toxic traditional Chinese medicinal materials⁽⁶⁷⁾.

(IV) The Limitation of Molecular Markers in Quality Control

There are several limitations. Firstly, it is not easy to extract DNA from some medicinal materials using general methods, particularly for those processed species. Secondly, although the differentiation of the geographical origins by molecular markers, such as the chloroplast *matK* gene sequence⁽⁶⁸⁾, ITS from nuclear rDNA⁽⁶⁹⁾, 18S rDNA gene⁽⁷⁰⁾ has been established from time to time, DNA markers may not correspond to the chemical profiles. Therefore, DNA markers together with the chemical fingerprint for quality control of CHM have been investigated. For example, three species of Rhizome *Curcuma* (Ezhu) including *C. wenyujin*, *C. phaeocalis*, and *C. kwangsiensis* have been used as medicinal materials. Chemical components such as curdione, curcumol, and germacrone in the essential oil are considered as the active constituents in *R. Curcumae*. The amount of these chemicals varies among samples from different species or

samples from the same species but from different regions of cultivation. Chemical fingerprints were generated from these species as the identification markers. At the same time, the 5S rDNA spacer domains of five *Curcuma* species, including the common adulterants of this herb, were sequenced. The chemical fingerprint together with the sequence data could serve as the marker for quality control of *Curcuma* species⁽⁵⁹⁾. To identify the origin of *Panax notoginseng* and its seven adulterants, and to analyze *P. notoginseng* in different localities, the nuclear 18S rDNA and chloroplast genes were sequenced. HPLC fingerprinting was also used to correlate the chemical composition and geographical distribution. This study concluded that DNA markers can be applied to authenticate the easily-confused species and can help to trace their geographical origins⁽⁷¹⁾. For the quality evaluation of *Pogostemon cablin* cultivated in Guangdong and Hainan, two sequences, 1.2 kb of plastid *matK* gene and 1.8 kb nuclear 18S rDNA gene, and two chemotypes (pogostone-type and Patchouliol-type in essential oil composition) were compared. The result showed that the sequence divergence in both *matK* and 18S rDNA genes among six samples of *P. cablin* were well correlated with their regions of cultivation and intraspecific chemotypes of essential oil compositions⁽⁷²⁾. Particularly, Chinese formulations noteworthy of multiple plant components make the identification more difficult, but it is not impossible. Testing for unknown contaminants is extremely difficult. These limitations are expected to be eliminated by advancement of molecular technology in the future.

CONCLUSIONS

Since the early paper concerning the differentiation of *P. quinquefolius* and *P. ginseng* by AP-PCR⁽²⁰⁾, many Chinese medicine materials have been authenticated by different molecular technologies (Table 2). Since more than 90% TCM uses either animal or plant as source material⁽⁷³⁾, molecular technologies may be an effective way to differentiate samples from different species or localities. Molecular markers have the advantages that they are least affected by age, environmental factors, and physiological conditions of the samples. These markers are not tissue-specific and thus can be detected at any stage of development. Moreover, a small amount of sample is sufficient for analysis and the physical form of the sample does not restrict the detection. These non-stringent requirements are particularly useful for some TCM that are expensive or in limited supply.

Urgent works for CHM quality control is to construct a comprehensive database of DNA fingerprints and DNA sequences for a broad spectrum of medicinal species. This comprehensive database containing voucher specimens, macro and microscopic data, chemical profiling and DNA fingerprinting information would clearly be beneficial for the authentication plant source, processing procedure, and




Figure 2. Microarray hybridization of individual herbal components in formulation A with Herba Dendrobii probe. Panels 1-9: spotted ITS1-5.8S-ITS2 sequences of the herbs 1-9 in formulation A. Panels P and N: spotted ITS1-5.8S-ITS2 DNA from formulation A with or without Herba Dendrobii. Row R: spotted ITS1-5.8S-ITS2 DNA of the five *Dendrobium* species listed in the Chinese Pharmacopoeia. The result of hybridization with ITS2 rDNA from *D. fimbriatum* (A) *D. nobile* (B) *D. officinale* (C) *D. loddigesii* (D) and *D. chrysanthum* (E) respectively were showed.

Table 2. List of Chinese medicines have been studied by different molecular technologies

Markers	TCM	Comments	Refs.
RAPD	<i>Glyrrhiza</i> species	Differentiation of four species	(74)
	<i>Zaocys dhumnades</i>	Identification of crude snake drugs	(75)
	<i>Anoectochilus</i>	Identification of two species	(76)
	<i>Atractylodes</i> plants	Revealed intraspecific variation	(77)
	<i>Astragalus</i> medicines	Differentiation of the two species	(78)
	<i>Rabdosis serra</i> plants	Authentication	(79)
	<i>Amomun villosum</i> species	Analysis of <i>A. villosum</i> and adulterants	(80)
	<i>Scutellaria</i> plants	Discrimination of the three species	(81)
	<i>Panax notoginseng</i>	Authentication of <i>P. notoginseng</i>	(82)
	Yu-ping-feng san	Identification of components	(83)
	<i>Aconitum</i> plants	Differentiation of <i>A. noveboracense</i> and <i>A. columbianum</i>	(84)
	<i>Ginkgo biloba</i>	Differentiation of the nine populations	(85)
	RFLP	<i>Atractylodes lancea</i>	Revealed intraspecific variation
<i>Panax</i> species		Differentiation of <i>P. ginseng</i> and <i>P. quinquefolius</i>	(87)
<i>Fritillaria pallidiflora</i>		Identification	(88)
rbcL	<i>Belamcanda chinensis</i>	Analysis of <i>B. chinensis</i> and related plants	(89)
trnK	<i>Curcuma</i> drugs	Authentication	(90)
	<i>Atractylodes</i> drugs	Authentication of derived crude drugs	(91)
	<i>Atractylodes</i> plants	Phylogenetic analysis	(92)
matK	<i>Panax vietnamensis</i>	Phylogenetic analysis	(93)
18S	<i>Panax notoginseng</i>	Analysis of homology	(94)
ITS	<i>Saussurea medusa</i>	Comparison on ITS sequences	(95)
	Herba Hedyotis Diffusae	Authentication	(96)
	<i>Hypericum</i> species	Genetic profiling	(97)
	<i>Zanthoxylum bungeanum</i> Maxim	Authentication of population and adulterants	(98)
5S	<i>Fritillaria</i> species	Molecular diversity	(99)
	Radix Astragali	Species identification	(100)
	<i>Ephedra</i> plants	Phylogenetic analysis	(101)
12S	Snake gallbladder	Identification	(102)
Cyt b	<i>Oviductus ranae</i>	Authentication of original animals	(103)

for providing consumers with a safe product. The molecular detection technologies therefore undoubtedly contribute to the research and development of herbal drugs.

ACKNOWLEDGEMENTS

This study was partially supported by a research grant (Project Code 200511159007) from the University of Hong Kong.

REFERENCES

- Bai, D. L. 1993. Traditional Chinese medicines and new drug development. *Pure Appl. Chem.* 65: 1103-1112.
- Shaw, P. C., Ngan, F. N., But, P. P. H. and Wang, J. 2002. Molecular markers in Chinese medicinal materials. In "Authentication of Chinese medicinal material by DNA technology". 1st ed. p. 1. Shaw, P. C., Wang, J. and But, P. P. H. eds. World Scientific Publishing. Singapore.
- Hon, C. C., Chow, Y. C., Zeng, F. Y. and Leung, F. C. C. 2003. Genetic authentication of ginseng and other traditional Chinese medicine. *Acta Pharmacol. Sin.* 24: 841-846.
- The State Pharmacopoeia Committee of the P. R. China. 2000. Dangsheng. In "Pharmacopoeia of P. R. China" (English Edition 2000). Volume I. p. 165. Chemical Industry Press. Beijing, China.
- Herbasin Chinese herb database. <http://www.herbasin.com/database/dangshen.htm>
- Li, J. and Xiao, X. 1995. An investigation on

- medicinal plant resources of *Dendrobium* in Sichuan province. *Chin. J. Chin. Mater. Med.* 20: 7-12.
7. Namba, T. and Lin, C. C. 1981. Pharmacognostical studies on the crude drugs of Orchidaceae from Taiwan (IV) on Chioh-hak. *Jpn. J. Pharmacog.* 35: 221-232.
 8. Zhang, J. H., Wang, Z. W., Zhang, X., Jin, J. Q. and Yang, F. L. 2000. Authentication of *Tribulus terrestris* L. by morphology, microscope and TLC. *Zhong Yao Cai.* 23: 675-677.
 9. Yin, J. X., Deng, X. H., Che, X. Y. and Zhang, L. H. 2005. Study on TLC identification of Fructus Xanthii. *W C J · P S.* 20: 067-069.
 10. He, Z. D., Qiao, C. F., Han, Q. B. Cheng, C. L., Xu, H. X, Jiang, R. W., But, P. P. H. and Shaw, P. C. 2005. Authentication and quantitative analysis on the chemical profile of cassia bark (cortex *cinnamomi*) by high-pressure liquid chromatography. *J. Agric. Food Chem.* 53: 2424-2428.
 11. Zhao, L., Huang, C., Shan, Z., Xiang, B. and Mei, L. 2005. Fingerprint analysis of *Psoralea corylifolia* L. by HPLC and LC-MS. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* 821: 67-74.
 12. Schaneberg, B. T., Crockett, S., Bedir, E. and Khan, I. A. 2003. The role of chemical fingerprinting: application to *Ephedra*. *Phytochemistry* 62: 911-918.
 13. Hu, P., Liang, Q. L., Luo, G. A., Zhao, Z. Z. and Jiang, Z. H. 2005. Multi-component HPLC fingerprinting of Radix Salviae Miltiorrhizae and its LC-MS-MS identification. *Chem. Pharm. Bull. (Tokyo).* 53: 677-683.
 14. Hao, G. M., Li, H. X., Zhao, C. J. and Zou, Y. Q. 2002. Determination of the content of atractylenoide III in the *Codonopsis pilosula* (Franch.) Nannf. by RP-HPLC. *J. Shenyang Pharm. Univ.* 19: 337-340.
 15. Zhu, H., Zhou, C. S, Liao, Y. K. and Bai, Y. Y. 2004. Pharmacognostic identification of *Cryptolepis buchanaei*. *China J. Chin. Mater. Med.* 29: 634-636.
 16. Ye, X., Yu, H. and Li, P. 2005. Analysis of Chinese drug beimu and its fake species with clustering analysis and FTIR spectra. *Zhong Yao Cai.* 28: 89-91.
 17. Ren, S. H., Li, Z. F., Zhao, Y., Chen, X. Q., Fu, L. N. and Xiao, C. P. 2004. Analysis of chemical constituents of volatile oil in armeniaca mume sieb by gas chromatography-mass spectrometry. *J. Taishan Med. Coll.* 25: 643-645.
 18. Xiao, S. Y., Luo, G. A., Wang, Y. M., Yang, X. D. and Liang, Q. L. 2004. Identification of Panax notoginseng and its preparations by LC/MS. *Acta Pharm. Sin.* 39: 127-131.
 19. Hu, P., Liang, Q. L., Luo, G. A., Zhao, Z. Z. and Jiang, Z. H. 2005. Multi-component HPLC fingerprinting of Radix Salviae Miltiorrhizae and its LC-MS-MS identification. *Chem. Pharm. Bull. (Tokyo).* 53: 677-683.
 20. Cheung, K. S., Kwan, H. S., But, P. P. H. and Shaw, P. C. 1994. Pharmacognostical identification of American and Oriental ginseng roots by genomic fingerprinting using arbitrarily-primed polymerase chain reaction. *J. Ethnopharmacol.* 42: 67-69.
 21. Shaw, P. C. and But, P. P. H. 1995. Authentication of *Panax* species and their adulterants by random-primed polymerase chain reaction. *Planta Med.* 61: 393-492.
 22. Zhang, Y. B., Leung, H. W., Yeung, H. W. and Wong, R. N. 2001. Differentiation of *Lycium barbarum* from its related Lycium species using random amplified polymorphic DNA. *Planta Med.* 67: 379-381.
 23. Um, J. Y., Chung, H. S., Kim, M. S., Na, H. J., Kwon, H. J., Kim, J. J., Lee, K. M., Lee, S. J., Lim, J. P., Do, K. R., Hwang, W. J., Lyu, Y. S., An, N. H. and Kim, H. M. 2001. Molecular authentication of *Panax ginseng* species by RAPD analysis and PCR-RFLP. *Biol. Pharm. Bull.* 24: 872-875.
 24. Cao, H., But, P. P. and Shaw, P. C. 1996. Authentication of the Chinese drug "ku-di-dan" (herba elephantopi) and its substitutes using random-primed polymerase chain reaction (PCR). *Acta Pharm. Sin.* 31: 543-553.
 25. Cao, H., But, P. P. and Shaw, P. C. 1997. Identification of herba taraxaci and its adulterants in Hong Kong market by DNA fingerprinting with random primed PCR. *Chin. J. Chin. Mater. Med.* 22: 197-200, 253.
 26. Fernandez, A., Garcia, T., Gonzalez, I., Asensio, L., Rodriguez, M. A., Hernandez, P. E. and Martin, R. 2002. Polymerase chain reaction-restriction fragment length polymorphism analysis of a 16S rDNA gene fragment for authentication of four clam species. *J. Food Prot.* 65: 692-695.
 27. Kaundun, S. S. and Matsumoto, S. 2003. Identification of processed Japanese green tea based on polymorphisms generated by STS-RFLP analysis. *J. Agric. Food Chem.* 51: 1765-1770.
 28. Yang, D. Y., Fushimi, H., Cai, S. Q. and Komatsu, K. 2004. Polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) and amplification refractory mutation system (ARMS) analyses of medicinally used Rheum species and their application for identification of Rhei Rhizoma. *Biol. Pharm. Bull.* 27: 661-669.
 29. Ha, W. Y., Shaw, P. C., Liu, J., Yau, F. C. and Wang, J. 2002. Authentication of *Panax ginseng* and *Panax quinquefolius* using amplified fragment length polymorphism (AFLP) and directed amplification of mini-satellite region DNA (DAMD). *J. Agric. Food Chem.* 50: 1871-1875.
 30. Ha, W. Y., Yau, F. C., But, P. P., Wang, J. and Shaw, P. C. 2001. Direct amplification of length polymorphism analysis differentiates *Panax ginseng* from *P. quinquefolius*. *Planta Med.* 67: 587-589.
 31. Yau, F. C., Wong, K. L., Wang, J., But, P. P. and Shaw, P. C. 2002. Generation of a sequence characterized amplified region probe for authentication of *Crocodilian* species. *J. Exp. Zool.* 294: 382-386.
 32. Wang, J., Ha, W. Y., Ngan, F. N., But, P. P. and Shaw, P. C. 2001. Application of sequence characterized amplified region (SCAR) analysis to authenticate *Panax* species and their adulterants. *Planta Med.* 67:

- 781-783.
33. Hon, C. C., Chow, Y. C., Zeng, F. Y. and Leung, F. C. 2003. Genetic authentication of ginseng and other traditional Chinese medicine. *Acta Pharmacol. Sin.* 24: 841-846.
 34. Zhang, Y. B., Ngan, F. N., Wang, Z. T., Wang, J., But, P. P. H. and Shaw, P. C. 1999. Differentiation of *Codonopsis pilosula* using random amplified polymorphic DNA. *Planta Med.* 65: 57-60.
 35. Li, H. L., Yang, Z., Du, H. X. and Liu, D. G. 2004. RAPD analysis of *Codonopsis lanceolata* in Jilin province. *J. Jilin. Agric. Univ.* 16: 66-69.
 36. Ding, G., Ding, X. Y., Shen, J., Tang, F., Liu, D. Y., He, J., Li, X. X and Chu, B. H. 2005. Genetic diversity and molecular authentication of wild populations of *Dendrobium officinale* by RAPD. *Acta Parma. Sin.* 40: 1028-1032.
 37. Tanaka, H., Fukuda, N. and Shoyama, Y. 2006. Identification and differentiation of *Panax* species using ELISA, RAPD and Eastern blotting. *Phytochem. Anal.* 17: 46-55.
 38. Wang, C. Z., Li, P., Ding, J. Y., Jin, G. Q. and Yuan, C. S. 2005. Identification of *Fritillaria pallidiflora* using diagnostic PCR and PCR-RFLP based on nuclear ribosomal DNA internal transcribed spacer sequences. *Planta Med.* 71: 384-386.
 39. Passinho-Soares, H., Felix, D., Kaplan, M. A., Margis-Pinheiro, M. and Margis, R. Authentication of medicinal plant botanical identity by amplified fragmented length polymorphism dominant DNA marker: Inferences from the *plectranthus* Genus. *Planta Med.* 2006 Jun 22; [Epub ahead of print].
 40. Hillis, D. M., Larson, A., Davis, S. K. and Zimmer, E. A. 1990. Nucleic acids III: sequencing. In "Molecular Systematics". pp. 318-370. Hillis, D. M. and Moritz, C. eds. Sunderland, MA: Sinauer Associates.
 41. Zhang, Y. B. 2000. Molecular approach to the authentication of *Lycium barbarum* and its related species. MPhil thesis, Hong Kong Baptist University, School of Chinese Medicine.
 42. Wang, C. Z., Li, P., Ding, J. Y., Jin, G. Q. and Yuan, C. S. 2005. Identification of *Fritillaria pallidiflora* using diagnostic PCR and PCR-RFLP based on nuclear ribosomal DNA internal transcribed spacer sequences. *Planta Med.* 71: 384-386.
 43. Chen, Y. Q., Hu, B., Xu, F., Zhang, W., Zhou, H. and Qu, L. H. 2004. Genetic variation of *Cordyceps sinensis*, a fruit-body-producing entomopathogenic species from different geographical regions in China. *FEMS Microbiol. Lett.* 230: 153-158.
 44. Liu, J. Q., Chen, Z. D., Liao, Z. X. and Lu, A. M. 2001. A comparison of its sequences of the Tibetan medicine "zang yin chen"--*Swertia mussotti* and its adulterant species. *Acta Pharm. Sin.* 36: 67-70.
 45. Cui, X. M., Lo, C. K., Yip, K. L., Dong, T. T. and Tsim, K. W. 2003. Authentication of *Panax notoginseng* by 5S rDNA spacer domain and random amplified polymorphic DNA (RAPD) analysis. *Planta Med.* 69: 584-586.
 46. Sun, Y., Fung, K. P., Leung, P. C., Shi, D. and Shaw, P. C. 2004. Characterization of medicinal *Epimedium* species by 5S rRNA gene spacer sequencing. *Planta Med.* 70: 287-288.
 47. Sasaki, Y., Fushimi, H., Cao, H., Cai, S. Q. and Komatsu, K. 2002. Sequence analysis of Chinese and Japanese Curcuma drugs on the 18S rRNA gene and *trnK* gene and the application of amplification-refractory mutation system analysis for their authentication. *Biol. Pharm. Bull.* 25: 1593-1599.
 48. Wong, K. L., Wang, J., But, P. P. and Shaw, P. C. 2004. Application of cytochrome b DNA sequences for the authentication of endangered *snake* species. *Forensic. Sci. Int.* 139: 49-55.
 49. Yang, M., Zhang, D., Liu, J. and Zheng, J. 2001. A molecular marker that is specific to medicinal rhubarb based on chloroplast *trnL/trnF* sequences. *Planta Med.* 67: 784-786.
 50. Fu, R. Z., Wang, J., Zhang, Y. B., Wang, Z. T., But, P. P. H., Li, N. and Shaw, P. C. 1999. Differentiation of medicinal *Codonopsis* species from adulterants by polymerase chain reaction-restriction fragment length polymorphism. *Planta Med.* 65: 648-650.
 51. Cosner, M. E., Raubeson, L. A. and Jansen, R. K. 2004. Chloroplast DNA rearrangements in Campanulaceae: phylogenetic utility of highly rearranged genomes. *BMC Evol. Biol.* 4: 27.
 52. Lau, T. W., Shaw, P. C., Wang, J. and But, P. P. 2001. Authentication of medicinal *Dendrobium* species by the internal transcribed spacer of ribosomal DNA. *Planta Med.* 67: 456-460.
 53. Xu, H., Li, X. B., Wang, Z. T., Ding, X. Y., Xu, L. S. and Zhou, K. Y. 2001. rDNA its sequencing of Herba *Dendrobii* (Huangcao). *Acta Pharm. Sin.* 36: 777-783.
 54. Ding, X. Y., Wang, Z. T., Xu, H., Xu, L. S., Zhou, K. Y. and Shi, G. X. 2002a. Study in sequence difference and SNP phenomenon of rDNA ITS region in F type population of *Dendrobium officinale*. *Chin. J. Chin. Mater. Med.* 27: 85-89.
 55. Ding, X., Xu, L., Wang, Z., Zhou, K., Xu, H. and Wang, Y. 2002b. Authentication of stems of *D. officinale* by rDNA ITS region sequences. *Planta Med.* 68: 191-192.
 56. Ding, X. Y., Wang, Z. T., Xu, H., Xu, L. S. and Zhou, K. Y. 2002c. Molecular authentication of *Dendrobium chrysanthum*. *Chin. J. Chin. Mater. Med.* 27: 407-411.
 57. Zhang, Y. B. 2004. DNA microarray for authentication of medicinal *Dendrobium* species. PhD thesis. The Chinese University of Hong Kong, Department of Biochemistry.
 58. Zhao, K. J., Dong, T. T., Cui, X. M., Tu, P. F. and Tsim, K. W. 2003. Genetic distinction of radix adenophorae from its adulterants by the DNA sequence of 5S-rRNA spacer domains. *Am. J. Chin. Med.* 31: 919-26.

59. Xia, Q., Zhao, K. J., Huang, Z. G., Zhang, P., Dong, T. T., Li, S. P. and Tsim, K. W. 2005. Molecular genetic and chemical assessment of *Rhizoma Curcumae* in China. *J. Agric. Food Chem.* 53: 6019-26.
60. Sasaki, Y., Fushimi, H., Cao, H., Cai, S. Q. and Komatsu, K. 2002. Sequence analysis of Chinese and Japanese *Curcuma* drugs on the 18S rRNA gene and *trnK* gene and the application of amplification-refractory mutation system analysis for their authentication. *Biol. Pharm. Bull.* 25: 1593-1599.
61. Zhu, S., Fushimi, H., Cai, S. and Komatsu, K. 2003. Phylogenetic relationship in the genus *Panax*: inferred from chloroplast *trnK* gene and nuclear 18S rRNA gene sequences. *Planta Med.* 69: 647-653.
62. Zhou, C., Zhou, K. and Zhang, S. 2004. Molecular authentication of the animal crude drug Sailonggu (bone of *Myospalax baileyi*). *Biol. Pharm. Bull.* 27: 1850-1858.
63. Mizukami, H., Okabe, Y., Kohda, H. and Hiraoka, N. 2000. Identification of the crude drug *atractylodes* rhizome (Byaku-jutsu) and *atractylodes lancea* rhizome (So-jutsu) using chloroplast *TrnK* sequence as a molecular marker. *Biol. Pharm. Bull.* 23: 589-594.
64. Guo, Y., Tsuruga, A., Yamaguchi, S., Oba, K., Iwai, K., Sekita, S. and Mizukami, H. 2006. Sequence analysis of chloroplast chlB gene of medicinal *Ephedra* species and its application to authentication of *Ephedra* Herb. *Biol. Pharm. Bull.* 29: 1207-1211.
65. Zhang, Y. B., Wang, J., Wang, Z. T., But, P. P. and Shaw, P. C. 2003. DNA microarray for identification of the herb of *Dendrobium* species from Chinese medicinal formulations. *Planta Med.* 69: 1172-1174.
66. Qin, J., Leung, F. C., Fung, Y., Zhu, D. and Lin, B. 2005. Rapid authentication of ginseng species using microchip electrophoresis with laser-induced fluorescence detection. *Anal. Bioanal. Chem.* 381: 812-819.
67. Carles, M., Cheung, M. K., Moganti, S., Dong, T. T., Tsim, K. W., Ip, N. Y. and Sucher, N. J. 2005. A DNA microarray for the authentication of toxic traditional Chinese medicinal plants. *Planta Med.* 71: 580-584.
68. Cao, H., Cai, J. N., Liu Y. P., Wang, Z. T. and Xu, L. S. 2001. Correlative analysis between geographical distribution and nucleotide sequence of chloroplast *matK* gene of *Cnidium monnieri* fruit in China. *Chin. Pharm. J.* 36: 373-376.
69. Liu, Y. P., Cao, H., Han, G. R., Fushimi, H. and Komatsu, K. 2002. *matK* and its nucleotide sequencing of crude drug chuanxiong and phylogenetic relationship between their species from China and Japan. *Acta Pharma. Sin.* 37: 63-68.
70. Fushimi, H., Komatsu, K., Namba, T. and Isobe M. 2000. Genetic heterogeneity of ribosomal RNA gene and *matK* gene in *Panax notoginseng*. *Planta Med.* 66: 659-661.
71. Zhang, Y., Huang, M. H., Bai, G. R., Yang, M. S. and Cao, H. 2005. Study on molecular and chemical identification of *Panax notoginseng* by 18S rRNA gene and *matK* gene sequencing and by HPLC fingerprinting. *Drug Evaluation* 2: 23-29.
72. Liu, Y. P., Luo, J. P., Feng, Y. F., Guo, X. L. and Cao, H. 2002. DNA Profiling of *pogostemon cablin* chemotypes differing in essential oil composition. *Acta Pharma. Sin.* 37: 304-308.
73. Yamazaki, M., Sato, A., Shimomura, K., Saito, K. and Murakoshi, I. 1994. Genetic relationships among *Glycyrrhiza* plants determined by RAPD and RFLP analyses. *Biol. Pharm. Bull.* 17: 1529-1531.
74. Xu, H., Wang, Z. T. and Hu, Z. B. 2003. Development and application of technology for DNA molecular identification in traditional Chinese medicine. *World Sci. Technol. Modernization Trad. Chin. Med. Materia Medica* 5: 24-30.
75. Wang, Y. Q. and Zhou, K. Y. 1997. A preliminary study on the identification of crude *snake* drugs by molecular genetic markers. *Acta Pharm. Sin.* 32: 384-387.
76. Hu, S. M., Zhang, Q. G., Zhou, H. T. and Ruan, R. B. 2000. Identification of two species of *Anoectochilus* Bl. (Jin Xian Lian) by RAPD. *Chin. Trad. Herb. Drug* 33: 949-950.
77. Kohjyouma, M., Nakajima, S., Namera, A., Shimizu, R., Mizukami, H. and Kohda, H. 1997. Random amplified polymorphic DNA analysis and variation of essential oil components of *Atractylodes* plants. *Biol. Pharm. Bull.* 20: 502-506.
78. Cheng, K. T., Su, B., Chen, C. T. and Lin, C. C. 2000. RAPD analysis of *Astragalus* medicines marketed in Taiwan. *Am. J. Chin. Med.* 28: 273-278.
79. Chen, L. J., Ju, L. H., Shi, S. H., Li, Z. H. and Lai, X. P. 1998. Application RAPD technology to authenticate *Rabdosin serra* plants. *Chin. J. Chin. Mater. Med.* 23: 328-382.
80. Wang, P., Huang, F., Zhou, L., Cao, L., Liang, S., Xu, H. and Liu, J. 2000. Analysis of *Amomun villosum* species and some adulterants of zingiberaceae by RAPD. *Zhong Yao Cai* 23: 71-74.
81. Hosokawa, K., Minami, M., Kawahara, K., Nakamura, I. and Shibata, T. 2000. Discrimination among three species of medicinal *Scutellaria* plants using RAPD markers. *Planta Med.* 66: 270-272.
82. Cui, X. M., Lo, C. K., Yip, K. L., Dong, T. T. and Tsim, K. W. 2003. Authentication of *Panax notoginseng* by 5S-rRNA spacer domain and random amplified polymorphic DNA (RAPD) analysis. *Planta Med.* 69: 584-586.
83. Cheng, K. T., Tsay, H. S., Chen, C. F. and Chou, T. W. 1998. Determination of the components in a Chinese prescription, yu-ping-feng san, by RAPD analysis. *Planta Med.* 64: 563-565.
84. Cole, C. T. and Kuchenreuther, M. A. 2001. Molecular markers reveal little genetic differentiation among *Aconitum noveboracense* and *A. columbianum* (Ranunculaceae) populations. *Am. J. Bot.* 88: 337-347.
85. Fan, X. X., Shen, L., Zhang, X., Chen, X. Y. and Fu, C. X. 2004. Assessing genetic diversity of *Ginkgo biloba*

- L. (Ginkgoaceae) populations from China by RAPD markers. *Biochem. Genet.* 42: 269-278.
86. Mizukami, H., Shimizu, R., Kohda, H., Kohjyouma, M., Kawanishi, F. and Hiraoka, N. 1996. Restriction fragment length polymorphisms of rDNA and variation of essential oil composition in *Atractylodes* plants. *Biol. Pharm. Bull.* 19: 577-580.
87. Ngan, F., Shaw, P., But, P. and Wang, J. 1999. Molecular authentication of *Panax* species. *Phytochemistry* 50: 787-791.
88. Wang, C. Z., Li, P., Ding, J. Y., Jin, G. Q. and Yuan, C. S. 2005. Identification of *Fritillaria pallidiflora* using diagnostic PCR and PCR-RFLP based on nuclear ribosomal DNA internal transcribed spacer sequences. *Planta Med.* 71: 384-386.
89. Qin, M. J., Huang, Y., Yang, G., Xu, L. S. and Zhou, K. Y. 2003. RbcL sequence analysis of *Belamcanda chinensis* and related medicinal plants of Iris. *Acta Pharm. Sin.* 38: 147-152.
90. Sasaki, Y., Fushimi, H., Cao, H., Cai, S. Q. and Komatsu, K. 2002. Sequence analysis of Chinese and Japanese Curcuma drugs on the 18S rRNA gene and *trnK* gene and the application of amplification-refractory mutation system analysis for their authentication. *Biol. Pharm. Bull.* 25: 1593-1599.
91. Mizukami, H., Okabe, Y., Kohda, H. and Hiraoka, N. 2000. Identification of the crude drug *atractylodes* rhizome (Byaku-jutsu) and *atractylodes lancea* rhizome (So-jutsu) using chloroplast *TrnK* sequence as a molecular marker. *Biol. Pharm. Bull.* 23: 589-594.
92. Mizukami, H., Shimizu, R., Kohjyouma, M., Kohda, H., Kawanishi, F. and Hiraoka, N. 1998. Phylogenetic analysis of *Atractylodes* plants based on chloroplast *trnK* sequence. *Biol. Pharm. Bull.* 21: 474-478.
93. Komatsu, K., Zhu, S., Fushimi, H., Qui, T. K., Cai, S. and Kadota, S. 2001. Phylogenetic analysis based on 18S rRNA gene and *matK* gene sequences of *Panax vietnamensis* and five related species. *Planta Med.* 67: 461-465.
94. Fushimi, H., Komatsu, K., Namba, T. and Isobe, M. 2000. Genetic heterogeneity of ribosomal RNA gene and *matK* gene in *Panax notoginseng*. *Planta Med.* 66: 659-661.
95. Liu, J. Q., Chen, Z. R. and Lu, A. M. 2001. Comparison on internal transcribed spacers (ITS) 3 sequences of Tibetan medicine *Saussurea medusa* and its easily confusable species. *Chin. Trad. Herb. Drug* 32: 443-445.
96. Liu, Z. and Hao, G. M. 2005. Authentication of the Herba Hedyotis Diffusae by rDNA ITS sequences. *Shannxi Zhong Yi* 26: 167-169.
97. Crockett, S. L., Douglas, A. W., Scheffler, B. E. and Khan, I. A. 2004. Genetic profiling of Hypericum (St. John's Wort) species by nuclear ribosomal ITS sequence analysis. *Planta Med.* 70: 929-935.
98. Shen, J., Ding, X. Y., Zhang, W. M., Bao, S. L., Chang, J. and Tang, F. 2005. Authentication of *Zanthoxylum bungeanum* Maxim population and adulterants by analysis of rDNA ITS sequences. *Acta Pharm. Sin.* 40: 80-86.
99. Cai, Z. H., Li, P., Dong, T. T. and Tsim, K. W. 1999. Molecular diversity of 5S-rRNA spacer domain in *Fritillaria* species revealed by PCR analysis. *Planta Med.* 65: 360-364.
100. Ma, X. Q., Duan, J. A., Zhu, D. Y., Dong, T. T. and Tsim, K. W. 2000. Species identification of Radix Astragali (Huangqi) by DNA sequence of its 5S-rRNA spacer domain. *Phytochemistry* 54: 363-368.
101. Long, C., Kakiuchi, N., Takahashi, A., Komatsu, K., Cai, S. and Mikage, M. 2004. Phylogenetic analysis of the DNA sequence of the non-coding region of nuclear ribosomal DNA and chloroplast of *Ephedra* plants in China. *Planta Med.* 70: 1080-1084.
102. Liu, X. H., Wang, Y. Q., Liu, Z. Q., Tong, Z. Z. and Zhou, K. Y. 2001. Identification of Chinese crude drug snake gallbladder by DNA molecular marker. *Acta Pharm. Sin.* 36: 229-232.
103. Xuegan, Y., Yiquan, W., Kaiya, Z. and Zhongquan, L. 2002. Authentication of oviductus ranae and its original animals using molecular marker. *Biol. Pharm. Bull.* 25: 1035-1039.