

In Vitro Evaluation of Antioxidant Activities and Inhibition of ACE Activity by Collagenase-Treated Hydrolysate Derived from Bullfrog Skin

YA-LING HUANG¹, SHU-HUI LI², TZU-YI CHUANG¹ AND CHAU-JEN CHOW^{1*}

¹ Department of Seafood Science, National Kaohsiung Marine University, 142, Hai-Chuan Road, Nan-Tzu, Kaohsiung 81157, Taiwan, R.O.C.

² Central of General Education, National Kaohsiung Marine University, 142, Hai-Chuan Road, Nan-Tzu, Kaohsiung 81157, Taiwan, R.O.C.

(Received: September 22, 2010; Accepted: May 6, 2011)

ABSTRACT

Enzymatic hydrolysates were obtained by hydrolysis of retorted skin gelatin hydrolysates (RSGH) from bullfrog (*Rana catesbeiana*) using three enzymes including collagenase, bromelain, and pancreatin for various times (0, 15, 30 and 60 min). As the hydrolysis time increased, degrees of hydrolysis (DH) of all three enzymatic hydrolysates increased. The molecular weight distribution of collagenase-hydrolyzed RSGH (RSGH-Cg) fraction was in the range of 189 - 686 Da (40.1 - 50.1%) and 686 - 6,511 Da (36.7 - 48.7%) while those of the other enzymatic-hydrolyzed RSGH fractions were mostly in the 686 - 6,511 Da (58.2 - 60.8%) range. All enzymatic hydrolysates had more potent antioxidant activity including 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity, ferrous ion chelating activity, and reducing power than the original RSGH. RSGH-Cg showed the highest activity. The lower IC₅₀ value of angiotensin I-converting enzyme (ACE) inhibitory activity of RSGH-Cg correlated with the degree of hydrolysis. This study suggested that collagenase treated bullfrog RSGH is a promising functional ingredient.

Key words: bullfrog, gelatin hydrolysate, enzymatic hydrolysis, antioxidant activity, ACE inhibitory activity

INTRODUCTION

The most abundant protein in the extracellular matrix of animal tissue is collagen. Gelatin is a series of polypeptide chains produced by denaturing collagen. Most commercial gelatins are derived from by-products of mammalian sources. In recent years, studies have investigated the properties of fish gelatin derived from cod⁽¹⁾, perch⁽²⁾, and tuna⁽³⁾. Other studies have reported that the properties of gelatin derived from fish skin, bone, and fin differed from those of mammalian gelatin due to differences in source and type of collagen^(4,5).

Enzymatic hydrolysis of proteins can produce hydrolysates with a wide range of applications as ingredients used in the nutritional, medical, and cosmetic industries. Several studies demonstrated that protein hydrolysates have marked antioxidant, blood pressure reducing and hypocholesterolemic activities⁽⁶⁻⁸⁾. Moreover, the gelatin hydrolysates from Alaskapollackskin⁽⁹⁾, seabreamscales⁽¹⁰⁾, and sea cucumber⁽¹¹⁾ have ACE inhibitory peptides that are antihypertensive.

Yang *et al.*⁽¹²⁾ reported that retorted skin gelatin hydrolysate from cobia (*Rachycentron canadum*) possesses several antioxidant effects, such as free-radical scavenging activity and inhibition of lipid peroxidation. From our observations, retorting procedure is the most important factor in the production of antioxidant hydrolysate from fish skin.

Many researchers have focused on the extraction of collagen derived from fish skin, bone, or scale as a replacement for mammalian collagen. The denaturing temperature of collagen from fish wastes is lower than that for porcine collagen⁽¹³⁾, but the lower melting point of the gelatin product makes it unsuitable for application as medical biomaterials. Li *et al.*⁽¹⁴⁾ pointed out that the acid-soluble collagen of bullfrog skin has higher thermal stability than mammalian collagens. Therefore, amphibian skin could be a promising source of collagen.

The family of bullfrog (*Rana catesbeiana*) belongs to the Ranidae family. Over 1,600 metric tonnes of bullfrog are cultured annually in southern Taiwan. Most bullfrog legs, after skin removal, are frozen and exported to other countries. Therefore, bullfrog skin, a by-product of processing, is a major fraction of frog wastes. A preliminary study showed

* Author for correspondence. Tel: +886-7-3617141 ext. 3607; Fax: +886-7-3658907; E-mail: cjchow@mail.nkmu.edu.tw

that fresh bullfrog skin contains approximately 25.3% protein that is acid-soluble collagen (~80% of total nitrogen content). The recovery of gelatin and its hydrolysate from fish skin collagen by thermal hydrolysis with retorting has been investigated in recent years. Bullfrog skin gelatin hydrolysate obtained by retorting treatment has been further converted to value-added enzymatic protein hydrolysates, which can be widely used as a potential bioactive substance. However, little information regarding the characteristics of gelatin hydrolysates derived from bullfrog skin has been reported. Therefore, the objective of this study was to characterize the enzymatic hydrolysates obtained from retorted bullfrog skin using enzymatic hydrolysis, by determining degree of hydrolysis and molecular weight distribution, to evaluate their antioxidant properties and angiotensin I-converting enzyme (ACE) inhibitory activity.

MATERIALS AND METHODS

I. Materials

The skin of bullfrog (*Rana catesbeiana*) was obtained from Ho-Wei Fishery Co., LTD, Pingtung County, Taiwan. After manual removal of limb tissue residues, the skin sample (30 ± 1.5 g per bullfrog) was washed, cut into small pieces (1.5×1.5 cm), placed in polyethylene bags, and stored at -20°C until use. All reagents were analytical grade.

II. Preparation of Bullfrog Retorted Skin Gelatin Hydrolysate (RSGH)

The frozen bullfrog skin was thawed to 4°C with tap water and soaked in 0.1 N NaOH with a skin: solution ratio of 1 : 10 (w/v) at 4°C for 3 h with gentle stirring. The skin sample was pooled and washed with tap water until neutral pH reached. The cleaned skin was further immersed in 6% phosphoric acid with a skin: solution ratio of 1 : 4 (w/v) for 3 h, drained and then washed under running tap water to remove excess acid. To obtain the gelatin hydrolysate, the swollen bullfrog skin was added with distilled water in 1 : 2 (w/v) ratio and hydrolyzed by retorting in an autoclave (HA-300M, Hirayama, Japan) at 121°C for 120 min. After filtration with metal sieve, the pH value of the retorted solution was 1.8 - 2.0. The solution was then neutralized with saturated $\text{Ca}(\text{OH})_2$ to pH 7. The neutralized solution was filtered again through a Buchner funnel with Toyo No. 5A filter paper to remove insoluble precipitants. The filtrate was dried by lyophilization and the dry matter was referred as 'RSGH' and stored at -20°C until further analysis.

III. Enzymatic Hydrolysis of RSGH Treated with Proteases

For further digestion of RSGH, the enzymatic hydrolysis was carried out using various enzymes including collagenase (from *Clostridium histolyticum*, NO. C0130, Sigma, St. Louis, MO, USA), bromelain (No. 1.01651.0025, Merck,

Darmstadt, Germany), and pancreatin (No. P1500, Sigma) at optimal activity temperature (37°C for collagenase, 50°C for bromelain, and 50°C for pancreatin). Dry RSGH sample (1 g) was digested with one of each protease in 0.1 M sodium phosphate buffer (pH 7.0) at an enzyme: hydrolysate ratio of 1 : 100 (w/w) and then hydrolyzed at optimal temperature for 0, 15, 30, and 60 min in a shaker-water bath with continuous stirring. The reaction was terminated in a boiling water bath for 10 min to inactivate enzyme activity. After centrifugation at $5,000 \times g$ for 10 min at 4°C , the supernatant was lyophilized and then stored at -20°C until use. The resultant product obtained from enzyme-treated RSGH was named RSGH-Bm (bromelain-treated), RSGH-Pc (pancreatin-treated), and RSGH-Cg (collagenase-treated), respectively.

IV. Determination of Degree of Hydrolysis (DH)

Following the method of Church *et al.*⁽¹⁵⁾, the degree of hydrolysis was determined with spectrophotometric assay using *o*-phthaldialdehyde (OPA). An aliquot (50 μL) was added to 1 mL OPA reagent. After mixing, the mixture was measured spectrophotometrically at 340 nm. Gelatin extracted at 50°C for 4 h from bullfrog skin was used as the control. The calibration curve is drawn using a series of concentrations of Leu-Gly as a reference, which is converted to the number of peptide bond. Degree of hydrolysis was defined as percentage ratio of the number of peptide bonds released after enzymatic hydrolysis of RSGH to the total number of peptide bonds of gelatin extracted at 50°C and was calculated as follows:

$$\text{DH (\%)} = \frac{\text{N}_h - \text{N}_g}{\text{N}_g} \times 100$$

where N_h was number of cleaved peptide bonds after enzymatic hydrolysis of RSGH and N_g was total number of peptide bonds of gelatin extracted at 50°C .

V. Molecular Weight Distribution

Molecular weight distributions of the hydrolysate samples were determined by the gel permeation chromatography using a Hitachi 2130 HPLC system equipped with Superdex peptide[®] 10/300 column (10×300 mm, Amersham Pharmacia Biotech, Uppsala, Sweden). Chromatography was carried out using 0.02 M phosphate buffer (pH 7.2) as mobile phase. An injection volume of 20 μL of each hydrolysate (2 mg protein/mL) and a flow rate of 0.5 mL/min and absorbance at 214 nm was used in the analysis. The molecular weight standard distributions were determined using the following standards including trypsin inhibitor (20 kDa, No. 202-09221, Wako, Osaka, Japan), cytochrome *C* (12.5 kDa, No. C2506, Sigma), aprotinin (6,511 Da, No. 52682, Serva Electrophoresis GmbH, Heideberg, Germany), pepstatin A (686 Da, No. 616370, Calbiochem, La Jolla, CA), Gly-Gly-Gly (189 Da, No. G1377, Sigma), and Gly (75 Da, No. 17-1323-01, Amersham Pharmacia Biotech). For quantitative determinations of molecular weight distribution (%), peak

area in the HPLC chromatograms was integrated in reference to the peak area of the standard proteins, using Hitachi EZChrom Elite 3.1 chromatography data system software.

VI. Scavenging Effect on 2,2-Diphenyl-1-picrylhydrazyl (DPPH) Radical

DPPH radical scavenging activity was determined by the method of Shimada *et al.*⁽¹⁶⁾ with slight modifications. Briefly, 0.1 mM solution of DPPH in 95% ethanol was prepared and 1 mL of this solution was added to 1 mL hydrolysate sample (6 mg/mL). The mixture was shaken vigorously and allowed to stand at room temperature for 30 min. Then the absorbance of resulting solution was measured spectrophotometrically at 517 nm. Lower absorbance of the reaction mixture indicates higher free radical scavenging activity. The percentage of DPPH scavenging effect was calculated using the following equation:

$$\text{DPPH scavenging effect (\%)} = 100 - \left(\frac{A_0 - A_1}{A_0} \right) \times 100$$

where A_0 was the absorbance of the control reaction and A_1 was the absorbance in the presence of the standard or hydrolysate sample.

VII. Determination of Reducing Power

The reducing power of hydrolysate sample was determined according to the method of Wu *et al.*⁽¹⁷⁾. An aliquot (1 mL) of each hydrolysate sample was mixed with 1 mL of 0.2 M phosphate buffer (pH 6.6) and 1 mL of 1% (w/v) potassium ferricyanide. The mixture was incubated at 20°C for 20 min and cooled rapidly. Then 1 mL of 10% (w/v) trichloroacetic acid was added to the mixture, which was then centrifuged at 1,000 ×g for 10 min. The upper layer of solution (2 mL) was mixed with distilled water (2 mL) and 0.1% (w/v) ferric chloride (0.5 mL) and allowed to stand in a dark room for 10 min. The absorbance of the resulting solution was measured spectrophotometrically at 700 nm. Higher absorbance of the reaction mixture indicated greater reducing power. L-Ascorbic acid (1,000 ppm) was used as a positive control.

VIII. Determination of Metal Ion Chelating Activity

The chelating of ferrous ions by the hydrolysate was measured using the method described by Wu *et al.*⁽¹⁷⁾ with slight modification. An aliquot (1 mL) of the hydrolysate sample was added with 2 mM FeCl_2 (100 μL). Subsequently, 200 μL of 5 mM ferrozine was added to this mixture. The reaction mixture was shaken vigorously and left standing at room temperature (24°C) for 10 min. The absorbance at 562 nm was then measured spectrophotometrically. The absorbance of 1,000 ppm EDTA was used as a positive control. Percentage of decrease in absorbance of the reaction mixture indicates increased metal ion chelating activity.

IX. Determination of ACE Inhibitory Activity

The ACE inhibitory activity was measured using the method described by Chushman and Cheung⁽¹⁸⁾ with slight modifications. The reaction mixture was prepared by adding 20 μL of 5 mM Hip-His-Leu into the 100 mM sodium borate buffer (pH 8.3) containing 300 mM NaCl as a substrate. An aliquot (80 μL) of hydrolysate treated with collagenase was mixed with the above reaction mixture and preincubated at 37°C for 5 min. Then, 20 μL of ACE (0.11 U/mL) was added and the mixture was incubated in a water bath at 37°C for 60 min and the reaction was stopped by adding 150 μL of 1.0 N HCl. Released hippuric acid was extracted by the addition of 1.7 μL ethyl acetate with continuous vortex mixing. After centrifuged at 1,600 ×g for 10 min, 1 mL of the upper layer was transferred into test tube and evaporated in vacuum. The resulting hippuric acid was redissolved in 1 mL distilled water and absorbance was measured spectrophotometrically at 280 nm. The concentration of hydrolysate caused a 50% reduction of ACE activity was expressed as the 50% inhibitory concentration (IC_{50}).

X. ACE Activity of Hydrolysates Following Digestion by Gastrointestinal Enzymes In Vitro

According to the methods of Wu and Ding⁽⁶⁾ and Chushman and Cheung⁽¹⁸⁾ with slight modifications, 1% (w/v) hydrolysate sample in 0.1 M KCl-HCl buffer (pH 2.0) was digested with pepsin in a water bath at 37°C for 4 h and placed in a boiling water bath for 15 min in order to terminate the enzymatic reaction. A portion of part of the mixture was subjected to the assay of ACE inhibitory activity and part of the mixture was neutralized with 2 N NaOH, followed by incubating with 2% (w/v) pancreatin in a 37°C water bath for 4 h. The reaction was stopped by heating in a 100°C water bath for 10 min. After centrifugation at 1,000 ×g for 10 min, the supernatant was used for ACE inhibitory activity determination. Inhibitory activity was expressed as the concentration of hydrolysate required to inhibit 50% of the ACE activity (IC_{50}).

XI. Statistical Analysis

All analyses were carried out in triplicate. Statistical analysis on the determinations was performed by one-way analysis of variance using the Statistical Analysis System (SAS). Values of $p < 0.05$ were considered statistically significant.

RESULTS AND DISCUSSION

In this study, the skin content of the whole bullfrog was about 6.0 g/100 g. After alkaline pretreatment and acid soaking, the gelatin hydrolysates obtained from the bullfrog skin were extracted with retorting in an autoclave, leading to a high nitrogen content of bullfrog RSGH. The nitrogen

content of the retorted filtrate reached approximately 50.1% of bullfrog skin (data not shown). Therefore, bullfrog skin can serve as a source of proteins. Yang *et al.*⁽¹²⁾ reported that the amount of nitrogen in cobia RSGH accounted for 46.9% of its nitrogen content in the skin, which is lower than that of bullfrog RSGH. Furthermore, degree of hydrolysis is used as a proteolysis monitoring parameter, and it is the most widely used indicator for comparison among different protein hydrolysates⁽¹⁹⁾. As shown in Figure 1, the extent of further degradation of RSGH by proteolytic enzymes was estimated by assessing the degree of hydrolysis. It was observed to be 83.1%, 74.3%, and 67.4%, respectively, for collagenase (37°C), bromelain (50°C), and pancreatin (50°C) after 15 min hydrolysis. Between 30 and 60 min of hydrolysis, no significant change in degree of hydrolysis of bullfrog RSGH was observed with all three proteases used. Among these hydrolysates, the degree of hydrolysis of RSGH treated with collagenase was the highest after 60 min hydrolysis. It is possible that collagenase contributed to a higher degree of hydrolysis due to its specific enzymatic action. During the retorting process, gelatin hydrolysate resulting from thermal denaturation produced from cobia skin was possibly attributed to the collagen solubility⁽¹²⁾. Moreover, the difference of degree of hydrolysis between the gelatin hydrolysates obtained by different proteases may be due to the extent of specific peptide bonds available for enzyme action^(19,20). Therefore, it was speculated that the degree of hydrolysis of gelatin hydrolysates was affected by the kind of raw material, as well as its pre-treatment (e.g. enzyme type) in the process.

The peak area of HPLC gel chromatography was evaluated for the molecular weight distribution of bullfrog RSGH during enzymatic hydrolysis. As shown in Table 1, bullfrog skin gelatin hydrolysates before any protease digestion gave rise to the dominant fraction of the smaller molecular weight, with 686 - 6,511 Da, which accounted for 54.0% of the total

hydrolysates. After 15 min of hydrolysis, all enzyme-treated RSGH had a size distribution of 686 - 6,511 Da. Moreover, the original RSGH fraction with a molecular weight larger than 6,511 Da was 26.5% of the hydrolysates. As hydrolysis time was increased to 60 min, the percentage of the RSGH-Cg, RSGH-Bm, and RSGH-Pc having molecular weights of over 6,511 Da significantly ($p < 0.05$) decreased to 0.9, 15.8, and 19.7%, respectively. Hence, the collagenase hydrolysis of retorted gelatin hydrolysates yielded more medium and small peptides. On the other hand, those having molecular weights below 686 Da were increased by treatment of collagenase (46.0 - 62.4%), bromelain (23.6 - 25.3%), and pancreatin (18.5 - 19.7%) from 15 min to 60 min hydrolysis.




Figure 1. Degree of hydrolysis of bullfrog RSGH by adding various proteases. With the same hydrolysis time, values (means \pm SD, $n = 3$) with different letters (a-a, g-i, p-r, w-y) are significantly different (Duncan, $p < 0.05$).

Table 1. The distribution of molecular weight fractions in the bullfrog RSGH treated with various proteases

Hydrolysate	Hydrolysis time (min)	Molecular weight ^a distribution (%) ^b				
		> 20,000	6,511 - 20,000	686 - 6,511	189 - 686	< 189
RSGH		1.4	25.1	54.0	14.2	5.3
RSGH-Cg ^c	15	0.0	3.2	48.7	40.1	5.9
	30	0.0	2.0	43.2	44.4	10.4
	60	0.0	0.9	36.7	50.1	12.3
RSGH-Bm ^d	15	0.2	18.0	58.2	17.8	5.8
	30	0.1	17.5	58.7	18.0	5.7
	60	0.1	15.7	58.8	19.4	5.9
RSGH-Pc ^e	15	0.3	20.8	60.4	13.1	5.4
	30	0.3	20.2	60.6	13.5	5.4
	60	0.3	19.4	60.8	14.1	5.6

^aThe molecular weight (Da) of all hydrolysates was evaluated using standard proteins as described in the Materials and Methods.

^bFor determination of molecular weight distribution (%), peak area in the HPLC chromatograms was integrated in reference to the peak area of the standard proteins. Each value represents the mean of triplicate determinations.

^{cde}RSGH-Cg, RSGH-Bm, and RSGH-Pc was obtained from RSGH treated with collagenase, bromelain, and pancreatin, respectively.

As a result, collagenase is capable of producing smaller peptides and peptides ranging from 189 to 686 Da account for almost half of the proteins in the collagenase-treated RSGH. A similar trend in molecular weight pattern was found in all enzyme-treated hydrolysate at hydrolysis times of 15 min to 60 min, but collagenase was the most effective protease of the enzymes tested. For all enzymes used in our study, the amount of small peptides obtained by enzymatic hydrolysis was apparently achievable within 60 min. Taken together, the molecular weights of each hydrolysate decreased distinctively as the peaks of high molecular weight fractions were reduced. Radha *et al.*⁽²¹⁾ studied the most evident change observed in the hydrolysate of oilseed protein in comparison to the untreated protein showed a reduction in the molecular weight of hydrolysate due to the presence of protease action. The production of enzymatic protein hydrolysates has smaller molecular sizes and less secondary structures than those of the original protein.

Enzymatic digestion has been extensively employed to improve the antioxidant properties of proteins. Gelatin hydrolysates produced by different proteolytic enzymes might have different molecular properties, leading to the varying characteristics of these resultant hydrolysates. Therefore, different methods (DPPH scavenging effect, metal ion chelating effect, and reducing power) need to be used to study their antioxidant properties. Generally, the effect of antioxidants on DPPH radical scavenging was due to their hydrogen donating ability. As depicted in Figure 2A, a significant ($p < 0.05$) decrease in the concentration of DPPH radical occurs with an increase concentration of the bullfrog RSGH and there is a good correlation between its concentration (2 - 8 mg/mL) and DPPH radical scavenging effects (20 - 72%). The DPPH scavenging effect of gelatin hydrolysate with further enzymatic digestion is shown in Figure 2B. The scavenging effect of the original hydrolysate at 6 mg/mL on DPPH radical was about 80%. As the hydrolysis time increased, there were no significant changes in DPPH scavenging activity for RSGH-Bm and RSGH-Pc. Among all three enzymatic hydrolysates, the treatment of RSGH with collagenase increased the maximal DPPH scavenging effect dependent on hydrolysis time (up to 87.0%). On the other hand, ascorbic acid exhibited 96.3% DPPH radical scavenging activity. RSGH-Cg may contain considerable amounts of small peptides that are electron donors and can react with free radicals to terminate the radical chain reaction. During hydrolysis, a wide variety of small peptides and free amino acids were generated which in turn might affect the resulting antioxidant activity^(22,23). Compared with a previous report, the DPPH scavenging activity (80 - 99%) of marine protein hydrolysates at a concentration of 40 mg/mL varied with different degree of hydrolysis⁽²⁰⁾. Bullfrog skin gelatin hydrolysate had a higher scavenging activity of DPPH radicals than those marine protein hydrolysates. As shown in Figure 3, the antioxidant activities of gelatin hydrolysates were evaluated using reducing power as an indicator. The result suggested that reducing power was markedly ($p < 0.05$) enhanced by ascorbic acid at the concentration of




Figure 2. (a) DPPH radical scavenging activity of different concentrations of bullfrog RSGH. (b) Effect of enzymatic hydrolysate derived from bullfrog RSGH and commercial antioxidant on DPPH radical scavenging activity. The concentration of all enzymatic hydrolysate used in all assays was 6 mg/mL, while that of ascorbic acid was 1,000 ppm. Values (means \pm SD, $n = 3$) with different letters (a-c) are statistically significant (Duncan, $p < 0.05$).




Figure 3. Reducing power of enzymatic hydrolysate derived from bullfrog RSGH and commercial antioxidant. The concentration of all enzymatic hydrolysates used in all assays was 6 mg/mL, while that of ascorbic acid was 1,000 ppm. Values (means \pm SD, $n = 3$) with different letters (a-d) are statistically significant (Duncan, $p < 0.05$).

1,000 ppm. By comparison, in the original RSGH sample, the reducing power (absorbance at 700 nm) of various enzymatic hydrolysates increased only slightly with increasing time of hydrolysis. With 15 min of hydrolysis, there was no significant change in the reducing power among all enzymatic hydrolysates. With an increased hydrolysis time to 60 min, RSGH-Cg exhibited more effective reducing power than the other two enzymatic hydrolysates. It appears that the reducing power of all gelatin hydrolysates depended on the enzyme used. Je *et al.*⁽²⁴⁾ demonstrated that protein hydrolysates of tuna liver showed a good reducing power, resulting from high contents of hydrogen-donating peptides from various enzymatic hydrolysates. Therefore, it is speculated that the RSGH-Cg having an effective antioxidant activity might be the result of increased exposure to a functional group such as hydroxyl and carboxyl with hydrogen donating abilities during thermal and proteolytic processes.

The ferrous ions chelating activity of all enzymatic hydrolysates and EDTA are shown in Figure 4. The percentage of chelating capacity of the original RSGH was found to be 50.5%, while that of EDTA reached almost 100%. Both RSGH-Cg and RSGH-Pc significantly ($p < 0.05$) increased the chelating ability with increased hydrolysis time. However, treatment of RSGH with bromelain for 15 - 60 min did not result in enhanced chelating activity after 15 min. For all enzymes used, gelatin hydrolysate with hydrolysis for 60 min exhibited higher metal ion chelating activity than that of RSGH. On the other hand, the RSGH-Bm, with 15-min hydrolysis, showed better chelating activity than the original RSGH and the other enzymatic hydrolysates. Therefore, an increase in the metal ion chelating activity was attributable to the hydrolytic action of certain enzymes. These results revealed that RSGH treated by various enzymes had a profound capacity for iron binding which might lead

to an action as a lipid peroxidation protector. The bullfrog skin gelatin hydrolysate shows a pronounced effect on the chelating metal ion and may retard the oxidation reaction.

ACE plays an important role in the rennin-angiotensin system (RAS), which can regulate both arterial blood pressure and salt and water balance. Captopril is an ACE inhibitor used for the treatment of hypertension. Some protein hydrolysates from whey, soy, milk, and globin⁽²⁵⁻²⁸⁾ contain small peptides that can significantly inhibit ACE activity. The ACE inhibition activity of all the hydrolysates of RSGH-Cg compared to that of Captopril is shown in Table 2. It is evident that the original RSGH had a very high IC_{50} value (0.193 mg/mL) of ACE inhibitory activity compared to those of enzymatic-hydrolysates at any hydrolysis time (0.024 - 0.073 mg/mL). The IC_{50} value of ACE inhibitory activity for collagenase-treated hydrolysate was significantly ($p < 0.05$) lowered with increasing hydrolysis time and reached a minimum value of 0.024 mg/mL at 60 min of hydrolysis. It can also be seen that high ACE inhibitory activity of collagenase-hydrolyzed hydrolysates was found within the 15 - 60 min hydrolysis period, and its IC_{50} value was significantly decreased by 62.2 - 87.6%. Other reports on enzymatic hydrolysates obtained from shrimp and Alaska pollack skin showed that their IC_{50} for ACE inhibitory activities was in the range of 0.795 - 0.980 mg/mL^(9,23). These results reveal that the small peptides with ACE inhibitory action were contained in the hydrolysates of bullfrog RSGH and that the ACE inhibitory activity of collagenase-treated hydrolysates was higher than those of marine hydrolysates. In Figure 1 and Table 2, there is an apparent correlation ($r^2 = -0.99$) between degree of hydrolysis of RSGH-Cg and IC_{50} value of ACE inhibitory activity, indicating that increasing hydrolysis decreased the ACE IC_{50} value. Hence, it demonstrated that hydrolysis with collagenase could be critical for release of ACE inhibitory peptides from an inactive form within the bullfrog RSGH. Matsui *et al.*⁽²⁹⁾ demonstrated that ACE inhibitory activity of alkaline protease hydrolysate from sardine muscle markedly increased with increasing proteolysis. As reported by Fahmi *et al.*⁽¹⁰⁾, IC_{50} values (0.57 mg/mL) for hydrolysate of sea bream scales were obtained by the hydrolysis of alkaline protease, which is higher than our observation on the bullfrog RSGH treated with collagenase. Thus, collagenase was selected for effective digestion of bullfrog RSGH in our study, and the dependence of collagenase-treated hydrolysates on the hydrolysis time distinctly improved the ACE inhibitory effect.

In order to predict the antihypertensive effects *in vivo*, the collagenase-treated hydrolysate obtained from 60 min hydrolysis, accompanied with original RSGH, was further analyzed for their ACE inhibitory activity using gastrointestinal proteases *in vitro*. Gastric *in vitro* incubation provides a practical and easy process to measure the stability of the peptide under oral administration. Table 3 indicates that RSGH-Cg had a relatively low ACE inhibitory activity ($IC_{50} = 0.024$ mg/mL) in reference to the IC_{50} of Captopril (0.0015 μ g/mL). The addition of pepsin or pepsin + pancreatin to the original RSGH fraction resulted in changes of




Figure 4. Chelating activity of enzymatic hydrolysate derived from bullfrog RSGH and commercial chelating reagent. The concentration of all enzymatic hydrolysates used in all assays was 6 mg/mL, while that of EDTA was 1,000 ppm. Values (means \pm SD, $n = 3$) with different letters (a-d) are statistically significant (Duncan, $p < 0.05$).

Table 2. ACE inhibitory activity (IC₅₀) of bullfrog RSGH treated with collagenase at various hydrolysis times

Hydrolysis time (min)	IC ₅₀ ^a
0	0.193 mg/mL
15	0.073 mg/mL
30	0.031 mg/mL
60	0.024 mg/mL
Captopril ^b	0.0015 µg/mL

^aEach value represents the mean of triplicate determinations.

^bCaptopril was used as a positive control. The IC₅₀ value of Captopril was cited from study of Tsai *et al.*⁽³⁰⁾.

Table 3. ACE inhibitory activity (IC₅₀) of the bullfrog RSGH and its RSGH-Cg after digestion by gastrointestinal protease

Gastrointestinal protease	IC ₅₀ ^a		
	RSGH (mg/mL)	RSGH-Cg ^b (mg/mL)	Captopril ^c (µg/mL)
Control ^d	0.193	0.024	0.0015
Pepsin	0.156	0.007	
Pepsin + pancreatin	0.428	0.014	

^aEach value represents the mean of triplicate determinations.

^bRSGH-Cg was obtained from RSGH treated with collagenase for 60 min at 37°C.

^cCaptopril was used as a positive control. The IC₅₀ value of Captopril was cited from study of Tsai *et al.*⁽³⁰⁾.

^dControl refers to the hydrolysate samples without adding any proteases.

IC₅₀ value (0.156 and 0.428 mg/mL, respectively) relative to that of undigested RSGH (0.193 mg/mL). In contrast, the ACE inhibitory activity (IC₅₀ = 0.007 mg/mL by pepsin, and IC₅₀ = 0.014 mg/mL by pepsin + pancreatin) for hydrolysis of RSGH-Cg had the higher activity than the undigested hydrolysate (0.024 mg/mL). The results indicated that RSGH-Cg obtained in the present study might be resistant to digestion in the gastrointestinal tract and exhibit a marked ACE inhibitory activity. Our findings agree with the study of Tsai *et al.*⁽³⁰⁾, who reported previously that hard clam muscle hydrolysate had an ACE inhibitory effect after pepsin or pepsin + pancreatin treatments. It is obvious that collagenase-treated hydrolysate could retain the ACE inhibitory activity after digestion of gastrointestinal proteases when orally administered. Thus, the RSGH-Cg with ACE inhibitory activity has low susceptibility to hydrolysis by gastrointestinal proteases.

CONCLUSIONS

Bullfrog skin enzymatic hydrolysate was obtained by a retorting treatment followed by proteolytic enzyme hydrolysis. The digestion of the resulting hydrolysate by collagenase exhibited good antioxidant properties. We also

found that bullfrog RSGH-Cg significantly decreased an IC₅₀ value of ACE inhibitory activity, suggesting a potential antihypertensive activity.

REFERENCES

- Kołodziejaska, I., Kaczorowski, K., Piotrowska, B. and Sadowska, M. 2004. Modification of the properties of gelatin from skins of Baltic cod (*Gadus morhua*) with transglutaminase. *Food Chem.* 86: 203-209.
- Muyonga, J. H., Cole, C. G. B. and Duodu, K. G. 2004. Extraction and physico-chemical characterisation of Nile perch (*Lates niloticus*) skin and bone gelatin. *Food Hydrocoll.* 18: 581-592.
- Je, J. Y., Qian, Z. J., Byun, H. G. and Kim, S. K. 2007. Purification and characterization of an antioxidant peptide obtained from tuna backbone protein by enzymatic hydrolysis. *Process Biochem.* 42: 840-846.
- Cho, S. M., Kwak, K. S., Park, D. C., Gu, Y. S., Ji, C. I., Jang, D. H., Lee, Y. B. and Kim, S. B. 2004. Processing optimization and functional properties of gelatin from shark (*Isurus oxyrinchus*) cartilage. *Food Hydrocoll.* 18: 573-579.
- Cheow, C. S., Norizah, M. S., Kyaw, Z. Y. and Howell, N. K. 2007. Preparation and characterisation of gelatins from the skins of sin croaker (*Johnius dussumieri*) and shortfin scad (*Decapterus macrosoma*). *Food Chem.* 101: 386-391.
- Wu, J. and Ding, X. 2002. Characterization of inhibition and stability of soy-protein-derived angiotensin I-converting enzyme inhibitory peptides. *Food Res. Int.* 35: 367-375.
- Sakanaka, S., Tachibana, Y., Ishihara, N. and Juneja, L. R. 2004. Antioxidant activity of egg-yolk protein hydrolysates in a linoleic acid oxidation system. *Food Chem.* 86: 99-103.
- Zhong, F., Liu, J., Ma, J. and Shoemaker, C. F. 2007. Preparation of hypocholesterol peptides from soy protein and their hypocholesterolemic effect in mice. *Food Res. Int.* 40: 661-667.
- Byun, H. G. and Kim, S. K. 2001. Purification and characterization of angiotensin I converting enzyme (ACE) inhibitory peptides from Alaska pollack (*Theragra chalcogramma*) skin. *Process Biochem.* 36: 1155-1162.
- Fahmi, A., Morimura, S., Guo, H. C., Shigematsu, T., Kida, K. and Uemura, Y. 2004. Production of angiotensin I converting enzyme inhibitory peptides from sea bream scales. *Process Biochem.* 39: 1195-1200.
- Zhao, Y., Li, B., Liu, Z., Dong, S., Zhao, X. and Zeng, M. 2007. Antihypertensive effect and purification of an ACE inhibitory peptide from sea cucumber gelatin hydrolysate. *Process Biochem.* 42: 1586-1591.
- Yang, J. I., Ho, H. Y., Chu, Y. J. and Chow, C. J. 2008. Characteristic and antioxidant activity of retorted gelatin hydrolysates from cobia (*Rachycentron canadum*) skin. *Food Chem.* 110: 128-136.

13. Nagai, T. and Suzuki, N. 2000. Isolation of collagen from fish waste material—skin, bone and fins. *Food Chem.* 68: 277-281.
14. Li, H., Liu, B. L., Gao, L. Z. and Chen, H. L. 2004. Studies on bullfrog skin collagen. *Food Chem.* 84: 65-69.
15. Church, F. C., Swaisgood, H. E., Porter, D.H. and Catignani, G. L. 1983. Spectrophotometric assay using *o*-phthaldialdehyde for determination of proteolysis in milk and isolated milk proteins. *J. Dairy Sci.* 66: 1219-1227.
16. Shimada, K., Fujikawa, K., Yahara, K. and Nakamura, T. 1992. Antioxidative properties of xanthan on the antioxidation of soybean oil in cyclodextrin emulsion. *J. Agric. Food Chem.* 40: 945-948.
17. Wu, H. C., Pan, B. S., Chang, C. L. and Shiau, C. Y. 2005. Low molecular weight peptides as related to antioxidant properties of chicken essence. *J. Food Drug Anal.* 13: 176-183.
18. Chushman, D. W. and Cheung, H. S. 1971. Spectrophotometric assay and properties of the angiotensin-converting enzyme of rabbit lung. *Biochem. Pharmacol.* 20: 1637-1648.
19. Guerard, F., Dufosse, L., De La Broise, D. and Binet, A. 2001. Enzymatic hydrolysis of protein from yellowfin tuna (*Thunnus albacares*) wastes using alcalase. *J. Mol. Catal. B: Enzym.* 11: 1051-1059.
20. Klompong, V., Benjakul, S., Kantachote, D. and Shahidi, F. 2007. Antioxidative activity and functional properties of protein hydrolysate of yellow stripe trevally (*Selaroides leptolepis*) as influenced by the degree of hydrolysis and enzyme type. *Food Chem.* 102: 1317-1327.
21. Radha, C., Kumar, P. R. and Parakash, V. 2008. Preparation and characterization of a protein hydrolysate from an oilseed flour mixture. *Food Chem.* 106: 1166-1174.
22. Wu, H. C., Chen, H. M. and Shiau, C. Y. 2003. Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). *Food Res. Int.* 36: 949-957.
23. He, H., Chen, X., Sun, C., Zhang, Y. and Gao, P. 2006. Preparation and functional evaluation of oligopeptide-enriched hydrolysate from shrimp (*Acetes chinensis*) treated with crude protease from *Bacillus* sp. SM98011. *Bioresour. Technol.* 97: 385-390.
24. Je, J. Y., Lee, K. H., Lee, M. H. and Ahn, C. B. 2009. Antioxidant and antihypertensive protein hydrolysates produced from tuna liver by enzymatic hydrolysis. *Food Res. Int.* 42: 1266-1272.
25. Mullally, M. M., Meisel, H. and Fitzgerald, R. J. 1997. Angiotensin-I-converting enzyme inhibitory activities of gastric and pancreatic proteinase digests of whey proteins. *Int. Dairy J.* 7: 299-303.
26. Gibbs, B. F., Zougman, A., Masse, R. and Mulligan, C. 2004. Production and characterization of bioactive peptides from soy hydrolysate and soy-fermented food. *Food Res. Int.* 37: 123-131.
27. Yu, Y., Hu, J., Bai, X., Du, Y. and Lin, B. 2006. Preparation and function of oligopeptide-enriched hydrolysate from globin by pepsin. *Process Biochem.* 41: 1589-1593.
28. Otte, J., Shalaby, S. M., Zakora, M., Pripp, A. H. and EI-Shabrawy, S. A. 2007. Angiotensin-converting enzyme inhibitory activity of milk protein hydrolysates: Effect of substrate, enzyme and time of hydrolysis. *Int. Dairy J.* 17: 488-503.
29. Matsui, T., Matsufuji, H., Seki, E., Osajima, K., Nakashima, M., Osajima, Y. 1993. Inhibition of angiotensin I-converting enzyme by *Bacillus licheniformis* alkaline protease hydrolyzates derived from sardine muscle. *Biosci Biotech Biochem* 57: 922-925.
30. Tsai, J. S., Chen, J. L. and Pan, B. S. 2008. ACE-inhibitory peptides identified from the muscle protein hydrolysate of hard clam (*Meretrix lusoria*). *Process Biochem.* 43: 743-747.